

Efficiency and strategy-proofness in object assignment problems with multi-demand preferences*

Tomoya Kazumura[†] and Shigehiro Serizawa[‡]

February 23, 2016

Abstract

We consider the problem of allocating sets of objects to agents and collecting payments. Each agent has a preference relation over the set of pairs consisting of a set of objects and a payment. Preferences are not necessarily quasi-linear. Non-quasi-linear preferences describe environments where the wealth effect is non-negligible: the payment level changes agents' willingness to pay for swapping sets. We investigate the existence of *efficient* and *strategy-proof* rules. A preference relation is *unit-demand* if given a payment level, for each set of objects, the most preferred one in the set is at least as good as the set itself; it is *multi-demand* if given a payment level, when an agent receives an object, receiving some additional object(s) makes him better off. We show that if a domain contains enough variety of unit-demand preferences and at least one multi-demand preference relation, and if there are more agents than objects, then no rule satisfies *efficiency*, *strategy-proofness*, *individual rationality*, and *no subsidy for losers* on the domain.

Keywords. Strategy-proofness, efficiency, multi-demand preferences, unit-demand preferences, non-quasi-linear preferences, minimum price Walrasian rule

JEL Classification Numbers. D44, D71, D61, D82

*The authors are grateful to the associate editor and two anonymous referees for their many detailed and helpful comments. The preliminary version of this article was presented at the IDGP 2015 Workshop, the 2015 Conference on Economic Design, and the II MOMA Meeting. The authors thank participants at those conferences for their comments. They are also grateful to seminar participants at Shanghai University of Finance and Economics, University of California, Berkeley, and Waseda University for their comments. The authors specially thank William Thomson, Yuichiro Kamada and Ryan Tierney for their detailed comments. This research was supported by the Joint Usage/Research Center at ISER, Osaka University. The authors acknowledge the financial support from the Japan Society for the Promotion of Science (Kazumura, 14J05972; Serizawa, 15H03328).

[†]Graduate School of Economics, Osaka University 1-7 Machikaneyama, Toyonaka, Osaka 560-0043, Japan. Email: pge003kt@student.econ.osaka-u.ac.jp)

[‡]Institute of Social and Economic Research, Osaka University, 6-1, Mihogaoka, Ibaraki, Osaka 567-0047, Japan. E-mail: serizawa@iser.osaka-u.ac.jp

1 Introduction

We consider an object assignment problem with money. Each agent receives a (possibly empty) set of objects and, possibly, pays money for the set. He has a preference relation over the set of pairs consisting of a set of objects and a payment. An *allocation* specifies how the objects are allocated and how much each agent pays. An (*allocation*) *rule* is a mapping from a class of admissible preference profiles, which we call a “domain,” to the set of allocations. An allocation is *efficient* if, without reducing the total payment, no other allocation makes all agents at least as well off and at least one agent better off. A rule is *efficient* if it always selects an efficient allocation. A rule is *strategy-proof* if, for each agent, it is a weakly dominant strategy to report his true preferences. We investigate the existence of *efficient* and *strategy-proof* rules.

Our model can be viewed as a multi-object auction model. Much of the literature on auction theory assumes preferences to be “quasi-linear.” This means that the valuations over sets of objects are not affected by payment level. On the quasi-linear domain, the so-called “VCG rules” (Vickrey, 1961; Clarke, 1971; Groves, 1973) are *efficient* and *strategy-proof*, and they are the only rules satisfying these properties (Holmström, 1979).

As Marshall (1920) demonstrates, preferences are approximately quasi-linear if payments are sufficiently low. However, in important applications of auction theory such as spectrum license allocation, house allocation, etc., prices are often equal to or exceed agents’ annual revenues. Excessive payments for objects may impair an agent’s ability to purchase complements for an effective use of the objects, and thus may influence the benefit the agent derives from the objects. Another reason why preferences may not be quasi-linear is that an agent may need a loan to be able to pay high prices, and typically financial costs are nonlinear in borrowing.^{1,2}

Another common assumption is the “unit-demand” property.³ It says that given a payment level, for each set of objects, the most preferred one in the set is at least as good as the set itself. For unit-demand preferences, there exists the *minimum Walrasian equilibrium price*, which is lower than any other Walrasian equilibrium price. Thus, on the “unit-demand domain,” minimum price Walrasian (MPW) rules are well-defined. A *minimum price Walrasian (MPW) rule* always selects an allocation associated with the minimum price Walrasian equilibria for each preference profile. The MPW rules are *strategy-proof* on the unit-demand domain (Demange and Gale, 1985). It is straightforward to see that the MPW rules satisfy the following additional two properties: One is *individual rationality*, which says that each agent finds his assignment at least as desirable as getting no object and paying nothing. The other is *no subsidy for losers*, which says that the payment of an agent who receives no object is nonnegative. On the unit-demand domain, when there are more agents than objects, the MPW rules are the only rules satisfying *efficiency*, *strategy-proofness*, *individual rationality*, and *no subsidy for losers* (Morimoto and Serizawa, 2015).⁴

Although the unit-demand assumption is suitable in some important cases such as house allocation, etc., in many other cases, some agents may well wish to receive more than one

¹ See Saitoh and Serizawa (2008) for numerical examples.

² Ausubel and Milgrom (2002) also discuss the importance of the analysis under non-quasi-linear preferences. Also see Sakai (2008) and Baisa (2013) for more examples of non-quasi-linear preferences.

³ For example, see Andersson and Svensson (2014), Andersson et al. (2015), and Tierney (2015).

⁴Note that the unit-demand domain contains non-quasilinear preferences, and thus the Hölmström (1979) result does not apply

object, and indeed, many authors have analyzed such situations.⁵

Now, a natural question arises. On a domain that is neither quasi-linear nor unit-demand, do *efficient* and *strategy-proof* rules exist? This is the question we address. To state our result, we need an additional property of preferences. A preference relation satisfies the *multi-demand* property if given a payment level, when an agent receives an object, receiving some additional object(s) makes him better off. We show that when there are more agents than objects, on any domain that contains enough variety of unit-demand preferences and at least one multi-demand preference relation, no rule satisfies *efficiency*, *strategy-proofness*, *individual rationality*, and *no subsidy for losers*. In most impossibility results in the literature on strategy-proofness, the incompatibility of a list of properties of rules is established on the fixed domain.⁶ On the other hand, our result is more general in the sense that the incompatibility of our properties holds on any domain containing enough variety of unit-demand preferences and some multi-demand preferences.

This article is organized as follows. In Section 2, we introduce the model and basic definitions. In Section 3, we introduce the unit-demand model and the richness condition. In Section 4, we define the minimum price Walrasian rule. In Section 5, we state our result and show the sketch of the proof. Section 6 concludes. All the proofs appear in the Appendix.

2 The model and definitions

There are $n \geq 2$ agents and $m \geq 2$ objects. We denote the set of agents by $N \equiv \{1, \dots, n\}$ and the set of objects by $M \equiv \{1, \dots, m\}$. Let \mathcal{M} be the power set of M . With abuse of notation, for each $a \in M$, we may write a to mean $\{a\}$. Each agent receives a subset of M and pays some amount of money. Thus, the agents' common **consumption set** is $\mathcal{M} \times \mathbb{R}$ and a generic (**consumption**) **bundle** for agent i is a pair $z_i = (A_i, t_i) \in \mathcal{M} \times \mathbb{R}$. Let $\mathbf{0} \equiv (\emptyset, 0)$.

Each agent i has a complete and transitive preference relation R_i over $\mathcal{M} \times \mathbb{R}$. Let P_i and I_i be the strict and indifference relations associated with R_i . A typical class of preferences is denoted by \mathcal{R} . We call \mathcal{R}^n a **domain**. The following are standard conditions of preferences.

Money monotonicity: For each $A_i \in \mathcal{M}$ and each pair $t_i, t'_i \in \mathbb{R}$ with $t_i < t'_i$, $(A_i, t_i) P_i (A_i, t'_i)$.

First object monotonicity: For each $(\{a\}, t_i) \in \mathcal{M} \times \mathbb{R}$, $(\{a\}, t_i) P_i (\emptyset, t_i)$.

Possibility of compensation: For each $(A_i, t_i) \in \mathcal{M} \times \mathbb{R}$ and each $A'_i \in \mathcal{M}$, there are $t'_i, t''_i \in \mathbb{R}$ such that $(A_i, t_i) R_i (A'_i, t'_i)$ and $(A'_i, t''_i) R_i (A_i, t_i)$.

Continuity: For each $z_i \in \mathcal{M} \times \mathbb{R}$, the **upper contour set at z_i** , $UC_i(z_i) \equiv \{z'_i \in \mathcal{M} \times \mathbb{R} : z'_i R_i z_i\}$, and the **lower contour set at z_i** , $LC_i(z_i) \equiv \{z'_i \in \mathcal{M} \times \mathbb{R} : z_i R_i z'_i\}$, are both closed.

Free disposal: For each $(A_i, t_i) \in \mathcal{M} \times \mathbb{R}$ and each $A'_i \in \mathcal{M}$ with $A'_i \subseteq A_i$, $(A_i, t_i) R_i (A'_i, t_i)$.

Definition 1 A preference relation is **classical** if it satisfies money monotonicity, first object monotonicity, possibility of compensation, and continuity.

⁵ For example, see Gul and Stacchetti (1999, 2000), Bikhchandani and Ostroy (2002), Papai (2003), Ausubel (2004, 2006), Mishra and Parkes (2007), de Vries et al (2007), and Sun and Yang (2006, 2009, 2014).

⁶ For example, Gibbard (1973), Satterthwaite (1975), etc.

Let \mathcal{R}^C be the class of classical preferences. We call $(\mathcal{R}^C)^n$ the **classical domain**. Let \mathcal{R}_+^C be a class of classical preferences satisfying free disposal. Obviously, $\mathcal{R}_+^C \subsetneq \mathcal{R}^C$.

Lemma 1 holds for classical preferences. The proof is relegated to the Appendix.

Lemma 1 *Let $R_i \in \mathcal{R}^C$, $z_i \in \mathcal{M} \times \mathbb{R}$, and $A'_i \in \mathcal{M}$. There is $t_i^* \in \mathbb{R}$ such that $z_i I_i (A'_i, t_i^*)$.*

For each $R_i \in \mathcal{R}^C$, each $z_i \in \mathcal{M} \times \mathbb{R}$, and each $A_i \in \mathcal{M}$, let $V_i(A_i; z_i) \in \mathbb{R}$ be such that $(A_i, V_i(A_i; z_i)) I_i z_i$. Call $V_i(A_i; z_i)$ the **valuation of A_i at z_i for R_i** . By continuity, for each $R_i \in \mathcal{R}^C$, each $(A_i, t_i) \in \mathcal{M} \times \mathbb{R}$, and each $A'_i \in \mathcal{M}$, $V_i(A_i; (A_i, t_i))$ is continuous with respect to t_i . By money monotonicity, for each $R_i \in \mathcal{R}^C$ and each pair $(A_i, t_i), (A'_i, t'_i) \in \mathcal{M} \times \mathbb{R}$, $(A_i, t_i) R_i (A'_i, t'_i)$ if and only if $V_i(A'_i; (A_i, t_i)) \leq t'_i$.

Definition 2 A preference relation $R_i \in \mathcal{R}^C$ is **quasi-linear** if for each pair $(A_i, t_i), (A'_i, t'_i) \in \mathcal{M} \times \mathbb{R}$ and each $t''_i \in \mathbb{R}$, $(A_i, t_i) I_i (A'_i, t'_i)$ implies $(A_i, t_i + t''_i) I_i (A'_i, t'_i + t''_i)$.

Let \mathcal{R}^Q be the class of quasi-linear preferences. We call $(\mathcal{R}^Q)^n$ the **quasi-linear domain**. Obviously, $\mathcal{R}^Q \subsetneq \mathcal{R}^C$.

Remark 1 Let $R_i \in \mathcal{R}^Q$. Then,

- (i) there is a **valuation function** $v_i : \mathcal{M} \rightarrow \mathbb{R}_+$ such that $v_i(\emptyset) = 0$, and for each pair $(A_i, t_i), (A'_i, t'_i) \in \mathcal{M} \times \mathbb{R}$, $(A_i, t_i) R_i (A'_i, t'_i)$ if and only if $v_i(A'_i) - t'_i \leq v_i(A_i) - t_i$, and
- (ii) for each $(A_i, t_i) \in \mathcal{M} \times \mathbb{R}$ and each $A'_i \in \mathcal{M}$, $V_i(A'_i; (A_i, t_i)) - t_i = v_i(A'_i) - v_i(A_i)$.

Now we define important classes of preferences. The following property formalizes the notion that given a payment level, an agent desires to consume at most one object.

Definition 3 A preference relation $R_i \in \mathcal{R}^C$ satisfies the **unit-demand** property if for each $(A_i, t_i) \in \mathcal{M} \times \mathbb{R}$ with $|A_i| > 1$, there is $a \in A_i$ such that $(a, t_i) R_i (A_i, t_i)$.^{7,8}

The condition means that given a payment level, for each set of objects, the most preferred one in the set is at least as good as the set itself. Note that it is possible that when an agent with a unit-demand preferences receives an object and his payment is fixed, an additional object makes him better off. However, this occurs only when he prefers the additional object to the original one. Figure 1 illustrates a unit-demand preference relation.

***** FIGURE 1 (Unit-demand preference relation) ENTERS HERE *****

Let \mathcal{R}^U be the class of unit-demand preferences. We call $(\mathcal{R}^U)^n$ the **unit-demand domain**. Obviously, $\mathcal{R}^U \subsetneq \mathcal{R}^C$.

We also consider a property that formalizes the notion that given a payment level, an agent desires to consume several objects.

Definition 4 A preference relation $R_i \in \mathcal{R}^C$ satisfies the **multi-demand** property if for each $(\{a\}, t_i) \in \mathcal{M} \times \mathbb{R}$, there is $A_i \in \mathcal{M}$ such that $a \in A_i$ and $(A_i, t_i) P_i (\{a\}, t_i)$.

⁷ Given a set X , $|X|$ denotes the cardinality of X .

⁸ Gul and Stacchetti (1999) define the unit-demand property for quasi-linear preferences. In their model, a preference relation $R_i \in \mathcal{R}^Q$ satisfies the unit-demand property if for each $A_i \in \mathcal{M}$ with $|A_i| > 1$, $v_i(A_i) = \max_{a \in A_i} v_i(a)$.

The condition says that given a payment level, when an agent receives an object, receiving some additional object(s) makes him better off. Note that given a payment level, even if an agent with a multi-demand preferences receives a set consisting of several objects, he may find it indifferent to each object in the set. Figure 2 illustrates a multi-demand preference relation.

***** FIGURE 2 (Multi-demand preference relation) ENTERS HERE *****

Let \mathcal{R}^M be the class of multi-demand preferences. We call $(\mathcal{R}^M)^n$ the **multi-demand domain**. The following are examples of preferences satisfying the multi-demand property.

Example 1: k -object-demand preferences. Given $k \in \{1, \dots, m\}$, a preference relation $R_i \in \mathcal{R}^C$ satisfies the **k -object-demand** property if (i) for each $(A_i, t_i) \in \mathcal{M} \times \mathbb{R}$ with $|A_i| < k$, and each $a \in M \setminus A_i$, $(A_i \cup \{a\}, t_i) P_i (A_i, t_i)$, and (ii) for each $(A_i, t_i) \in \mathcal{M} \times \mathbb{R}$ with $|A_i| \geq k$, there is $A'_i \subseteq A_i$ with $|A'_i| = k$ such that $(A'_i, t_i) I_i (A_i, t_i)$.⁹ Clearly, for each $k \in \{2, \dots, m\}$, preferences satisfying the k -object-demand property satisfy the multi-demand property.

Example 2: Substitutes and complements. Suppose that the set of objects are divided into two non-empty sets K and L , and agent i with a preference relation R_i views objects a and b as substitutes if both a and b are in the same set, and as complements if a and b are in different sets. For example, objects in K can be pens and objects in L can be notebooks. Formally, R_i satisfies the following property: For each $A_i \in \mathcal{M}$ with $|A_i| > 1$ and each $t_i \in \mathbb{R}$, if $A_i \subseteq K$ or $A_i \subseteq L$, then there is $a \in A_i$ such that $(A_i, t_i) I_i (a, t_i)$, and otherwise, for each $a \in A_i$, $(A_i, t_i) P_i (a, t_i)$. Clearly, this preference relation R_i satisfies the multi-demand property.

Example 3: Object monotonic preferences with capacity. Consider the following situation: Cars are assigned to agents. Each agent owns a garage where he can park at most three cars, and if he has more than three cars, he needs to park some of them at outside the garage with some amount of fee. Then, an agent may have the following preferences R_i : for each $(A_i, t_i) \in \mathcal{M} \times \mathbb{R}$ and each $a \in M \setminus A_i$, if $|A_i| \leq 2$, then $(A_i \cup \{a\}, t_i) P_i (A_i, t_i)$, and otherwise, $(A_i, t_i) P_i (A_i \cup \{a\}, t_i)$. This preference relation satisfies the multi-demand property.

Some preferences in \mathcal{R}^C violate both of the unit-demand property and the multi-demand property.

Example 4: (Figure 3.) A preference relation violating the unit-demand property and the multi-demand property. Let $R_i \in \mathcal{R}^C$ be such that for each $a \in M$ and each $t_i \in \mathbb{R}$, $V_i(a; (\emptyset, t_i)) = t_i + 5$, and for each $A_i \in \mathcal{M}$ with $|A_i| > 1$, and each $t_i \in \mathbb{R}$,

$$V_i(A_i; (\emptyset, t_i)) = \begin{cases} t_i + 5 & \text{if } t_i \geq -5, \\ \frac{1}{2}(t_i + 5) & \text{otherwise.} \end{cases}$$

Then, for each pair $a, b \in M$ and each $t_i \in \mathbb{R}$ with $t_i < -5$, $V_i(\{a, b\}; (\emptyset, t_i)) = \frac{1}{2}(t_i + 5) > t_i + 5 = V_i(a; (\emptyset, t_i)) = V_i(b; (\emptyset, t_i))$, and thus, we have $(\{a, b\}, t_i + 5) P_i (a, t_i + 5) I_i (b, t_i + 5)$. Thus, R_i does not satisfy the unit-demand property. Moreover, for each $a \in M$, each $A_i \in \mathcal{M}$ with $a \in A_i$, and each $t_i \in \mathbb{R}$ with $t_i \geq -5$, $V_i(A_i; (\emptyset, t_i)) = t_i + 5 = V_i(a; (\emptyset, t_i))$, and thus, we have $(A_i, t_i + 5) I_i (a, t_i + 5)$. Thus, R_i does not satisfy the multi-demand property.

⁹ In Gul and Stacchetti (1999), this notion is called *k-satiation*

***** FIGURE 3 (R_i in Example 4) ENTERS HERE *****

An **object allocation** is an n -tuple $A \equiv (A_1, \dots, A_n) \in \mathcal{M}^n$ such that $A_i \cap A_j = \emptyset$ for each $i, j \in N$ with $i \neq j$. We denote the set of object allocations by \mathcal{A} . A **(feasible) allocation** is an n -tuple $z \equiv (z_1, \dots, z_n) \equiv ((A_1, t_1), \dots, (A_n, t_n)) \in (\mathcal{M} \times \mathbb{R})^n$ such that $(A_1, \dots, A_n) \in \mathcal{A}$. We denote the set of feasible allocations by Z . Given $z \in Z$, we denote the object allocation and the agents' payments at z by $A \equiv (A_1, \dots, A_n)$ and $t \equiv (t_1, \dots, t_n)$, respectively, and we also write $z = (A, t)$.

A **preference profile** is an n -tuple $R \equiv (R_1, \dots, R_n) \in \mathcal{R}^n$. Given $R \in \mathcal{R}^n$ and $i \in N$, let $R_{-i} \equiv (R_j)_{j \neq i}$.

An **allocation rule**, or simply a **rule** on \mathcal{R}^n is a function $f : \mathcal{R}^n \rightarrow Z$. Given a rule f and $R \in \mathcal{R}^n$, we denote the bundle assigned to agent i by $f_i(R)$ and we write $f_i(R) = (A_i(R), t_i(R))$.

Now, we introduce standard properties of rules. The efficiency notion here takes the planner's preferences into account and assume that he is only interested in his revenue. Formally, an allocation $z \equiv ((A_i, t_i))_{i \in N} \in Z$ is **(Pareto-)efficient** for $R \in \mathcal{R}^n$ if there is no feasible allocation $z' \equiv ((A'_i, t'_i))_{i \in N} \in Z$ such that (i) for each $i \in N$, $z'_i R_i z_i$, (ii) for some $j \in N$, $z'_j P_j z_j$, and (iii) $\sum_{i \in N} t'_i \geq \sum_{i \in N} t_i$.

The first property states that for each preference profile, a rule chooses an efficient allocation.

Efficiency: For each $R \in \mathcal{R}^n$, $f(R)$ is efficient for R .

Remark 2 By money monotonicity and Lemma 1, the efficiency of allocation z is equivalent to the property that there is no allocation $z' \equiv ((A'_i, t'_i))_{i \in N} \in Z$ such that

(i') for each $i \in N$, $z'_i I_i z_i$, and (ii') $\sum_{i \in N} t'_i > \sum_{i \in N} t_i$.

The second property states that no agent benefits from misrepresenting his preferences.

Strategy-proofness: For each $R \in \mathcal{R}^n$, each $i \in N$, and each $R'_i \in \mathcal{R}$, $f_i(R) R_i f_i(R'_i, R_{-i})$.

The third property states that an agent is never assigned a bundle that makes him worse off than he would be if he had received no object and paid nothing.

Individual rationality: For each $R \in \mathcal{R}^n$ and each $i \in N$, $f_i(R) R_i \mathbf{0}$.

The fourth property states that the payment of each agent is always nonnegative.

No subsidy: For each $R \in \mathcal{R}^n$ and each $i \in N$, $t_i(R) \geq 0$.

The final property is a weaker variant of the fourth: If an agent receives no object, his payment is nonnegative.

No subsidy for losers: For each $R \in \mathcal{R}^n$ and each $i \in N$, if $A_i(R) = \emptyset$, $t_i(R) \geq 0$.

3 Unit-demand model and rich domains

In our model, potentially each agent can receive several objects. However, some authors study a model in which no agent can receive more than one object owing to some reason, say by regulations, or by physical reasons.¹⁰ We call this model the *unit-demand model*, and refer

¹⁰ For example, Alkan and Gale (1990), Demange and Gale (1985), etc.

our model as the *multi-demand model*. Some important results are established in the unit-demand model, and they are related to our main result. Some of such results continue to hold in the multi-demand model when preferences are unit-demand, and others continue to hold only when domains include enough variety of unit-demand preferences. In this section, we introduce “richness” of domains in our model, which guarantees that a domain of the multi-demand model includes enough variety of unit-demand preferences.

In the unit-demand model, preferences are defined over $M \cup \{0\} \times \mathbb{R}$, where 0 means not receiving any object in M and is called *null object*. To distinguish classes of preferences in the multi-demand model and those in the unit-demand model, we denote a typical class of preferences in the unit-demand model by \mathcal{R} .

Money monotonicity, first object monotonicity, possibility of compensation, and continuity are defined in the unit-demand model in the same manner as defined in our model. Thus, in the unit-demand model, classical preferences are defined in the same manner.

Definition 5 A preference relation R_i over $M \cup \{0\} \times \mathbb{R}$ is **classical** if it satisfies money monotonicity, first object monotonicity, possibility of compensation, and continuity.

Let \mathcal{R}^C be the class of classical preferences in the unit-demand model.

In the unit-demand model, a feasible allocation is an n -tuple $z = ((x_i, t_i))_{i \in N} \in (M \cup \{0\} \times \mathbb{R})^n$ such that for each pair $i, j \in N$, $x_i = x_j$ implies $x_i = x_j = 0$. As we mentioned, in the unit-demand model, no agent can receive more than one object. Other notions such as rules, properties of rules, etc., are defined in the same manner as defined in the multi-demand model.

To define the richness, we introduce the following notions, which connect preferences in the multi-demand model to those in the unit-demand model.

Definition 6 A preference relation R_i in the multi-demand model **induces** a preference relation R'_i over $M \cup \{0\} \times \mathbb{R}$ if for each pair $(a, t_i), (b, t'_i) \in M \cup \{0\} \times \mathbb{R}$, $(a, t_i) R'_i (b, t'_i)$ if and only if $(A_i, t_i) R_i (A'_i, t'_i)$, where

$$A_i = \begin{cases} \{a\} & \text{if } a \in M, \\ \emptyset & \text{if } a = 0, \end{cases} \quad \text{and} \quad A'_i = \begin{cases} \{b\} & \text{if } b \in M, \\ \emptyset & \text{if } b = 0, \end{cases}$$

Definition 7 A class of preferences \mathcal{R} in the multi-demand model **induces** a class of preferences \mathcal{R}' over $M \cup \{0\} \times \mathbb{R}$ if (i) for each $R'_i \in \mathcal{R}'$, there is $R_i \in \mathcal{R}$ that induces R'_i , and (ii) for each $R_i \in \mathcal{R}$, there is $R'_i \in \mathcal{R}'$ that is induced by R_i .

Remark 3 Each of \mathcal{R}^U , \mathcal{R}_+^U , and $\mathcal{R}^U \setminus \mathcal{R}_+^U$ induces \mathcal{R}^C .

Now, we introduce the richness of domain, which guarantees that a domain of the multi-demand model includes enough variety of unit-demand preferences so that they induce the class of classical preferences in the unit-demand model.

Definition 8 A class of preferences \mathcal{R} is **rich** if $\mathcal{R} \cap \mathcal{R}^U$ induces \mathcal{R}^C .

By Remark 3, \mathcal{R}^U , \mathcal{R}_+^U , and $\mathcal{R}^U \setminus \mathcal{R}_+^U$ are rich.

4 Minimum price Walrasian rules

In this section we define the minimum price Walrasian rules and state several facts related to them.

Let $p \equiv (p^1, \dots, p^M) \in \mathbb{R}_+^m$ be a price vector. The **budget set at p** is defined as $B(p) \equiv \{(A_i, t_i) \in \mathcal{M} \times \mathbb{R} : t_i = \sum_{a \in A_i} p^a\}$. Given $R_i \in \mathcal{R}$, the **demand set at p for R_i** is defined as $D(R_i, p) \equiv \{z_i \in B(p) : \text{for each } z'_i \in B(p), z_i R_i z'_i\}$.

Lemma 2 *Let $R_i \in \mathcal{R}^U$ and $p \in \mathbb{R}_+^m$. (i) Suppose $p \in \mathbb{R}_{++}^m$. Then, for each $(A_i, t_i) \in D(R_i, p)$, $|A_i| \leq 1$. (ii) Let $A_i \in \mathcal{M}$ be such that $(A_i, \sum_{a \in A_i} p^a) R_i (A'_i, \sum_{a \in A'_i} p^a)$ for each $A'_i \in \mathcal{M}$ with $|A'_i| \leq 1$. Then, $(A_i, \sum_{a \in A_i} p^a) \in D(R_i, p)$.*

Definition 9 Let $R \in \mathcal{R}^n$. A pair $((A, t), p) \in Z \times \mathbb{R}_+^m$ is a **Walrasian equilibrium (WE)** for R if

W-i: for each $i \in N$, $(A_i, t_i) \in D(R_i, p)$, and

W-ii: for each $a \in M$, if $a \notin A_i$ for each $i \in N$, then, $p^a = 0$.

Condition W-i says that each agent receives a bundle that he demands. Condition W-ii says that an object's price is zero if it is not assigned to anyone. Given $R \in \mathcal{R}^n$, let $W(R)$ and $P(R)$ be the sets of Walrasian equilibria and prices for R , respectively.

Lemma 3 *Let $R \in (\mathcal{R}^U)^n$ and $p \in P(R)$. (i) If $n > m$, then $p^a > 0$ for each $a \in M$. (ii) There is $((A, t), p) \in W(R)$ such that $|A_i| \leq 1$ for each $i \in N$.*

Let $R \in (\mathcal{R}^U)^n$ and R' be preference profiles of the multi-demand and unit-demand models respectively such that for each $i \in N$, R'_i is induced by R_i . For each $p \in P(R)$, by (ii) of Lemma 3, there is an allocation $((\{a_i\}, t_i))_{i \in N}$ such that $((\{a_i\}, t_i))_{i \in N}, p) \in W(R)$, and thus, $((a_i, t_i))_{i \in N}, p)$ is a WE for R' . On the other hand, for each $p \in \mathbb{R}_+^m$, if p is a WE price vector for R' , then there is an allocation $((a_i, t_i))_{i \in N}$ such that $((a_i, t_i))_{i \in N}, p)$ is a WE for R' , and thus by (ii) of Lemma 2, $((\{a_i\}, t_i))_{i \in N}, p) \in W(R)$. Therefore, the set of WE price vectors for R coincides with the set of WE price vectors for R' .

In the unit-demand model, several results on Walrasian equilibrium are established. By the preceding argument, the same results continue to hold in our model for preferences satisfying the unit-demand property.

Fact 1 (Alkan and Gale, 1990) ^{11, 12} *For each $R \in (\mathcal{R}^U)^n$, a Walrasian equilibrium for R exists.*

Fact 2 (Demange and Gale, 1985) *For each $R \in (\mathcal{R}^U)^n$, there is a unique minimum Walrasian equilibrium price vector, i.e., a vector $p \in P(R)$ such that for each $p' \in P(R)$, $p \leq p'$.*¹³

¹¹ Precisely, Alkan and Gale (1990) show the non-emptiness of the core in a two-sided matching model. However, the two-sided model includes the unit-demand model, and in the unit-demand model, non-emptiness of the core is equivalent to the existence of a Walrasian equilibrium.

¹² Fact 1 is also shown by other authors. See, for example, Quinzi (1984), Gale (1984).

¹³ For each $p, p' \in \mathbb{R}^m$, $p \leq p'$ if and only if for each $i \in \{1, \dots, m\}$, $p^i \leq p'^i$.

A **minimum price Walrasian equilibrium (MPWE)** is a Walrasian equilibrium whose price is minimum. Given $R \in \mathcal{R}^n$, let $p_{\min}(R)$ be the **minimum Walrasian equilibrium price for R** , and $Z_{\min}^W(R)$ be the **set of Walrasian equilibrium allocations associated with $p_{\min}(R)$** . Although there might be several minimum price Walrasian equilibria, they are indifferent for each agent, *i.e.*, for each $R \in \mathcal{R}^n$, each pair $z, z' \in Z_{\min}^W(R)$, and each $i \in N$, $z_i I_i z'_i$.

Definition 10 A rule f on \mathcal{R}^n is a **minimum price Walrasian (MPW) rule** if for each $R \in \mathcal{R}^n$, $f(R) \in Z_{\min}^W(R)$.

It is easy to show that the MPW rules on $(\mathcal{R}^U)^n$ satisfy *efficiency, individual rationality, and no subsidy*. Demange and Gale (1985) show that the MPW rules are *strategy-proof* on the classical domain in the unit-demand model. Our arguments above allow us to convert each MPWE allocation in the unit-demand *model* into an MPWE for a unit-demand *profile* in the multi-demand model. Moreover, all the minimum price Walrasian equilibria are indifferent for each agent. Thus, the result by Demange and Gale (1985) implies that for each rich class of preferences $\mathcal{R} \subseteq \mathcal{R}^U$, the MPW rules on \mathcal{R}^n also satisfy *strategy-proofness* in multi-demand model.

Morimoto and Serizawa (2015) shows that in the unit-demand model, when $n > m$, only the MPW rules satisfy *efficiency, strategy-proofness, individual rationality, and no subsidy for losers* on $(\mathcal{R}^C)^n$. The following lemma states that in the multi-demand model, when $n > m$, *efficient* rules never assign more than one object to agents whose preferences satisfy the unit-demand property.

Lemma 4 (Single object assignment) *Let $n > m$. Let $\mathcal{R} \subseteq \mathcal{R}^C$ and f be an efficient rule on \mathcal{R}^n . Let $R \in \mathcal{R}^n$ and $i \in N$. If $R_i \in \mathcal{R}^U$, $|A_i(R)| \leq 1$.*

By Lemma 4, when $n > m$, for each rich class of preferences $\mathcal{R} \subseteq \mathcal{R}^U$, and each rule on \mathcal{R}^n satisfying *efficiency*, it always assigns each agent at most one object. Thus, when $n > m$, for each $\mathcal{R} \subseteq \mathcal{R}^U$ that is rich, and each rule on \mathcal{R}^n satisfying *efficiency, strategy-proofness, individual rationality, and no subsidy for losers*, there is a corresponding rule in the unit-demand model, and moreover, it is easy to see that the corresponding rule also satisfies the four properties. Thus, the result by Morimoto and Serizawa (2016) continues to hold in our model.

Fact 3 (Demange and Gale, 1985 for (i); Morimoto and Serizawa, 2015 for (ii)) *Let $\mathcal{R} \subseteq \mathcal{R}^U$. (i) The minimum price Walrasian rules on \mathcal{R}^n satisfy efficiency, strategy-proofness, individual rationality and no subsidy. (ii) Let $n > m$, and \mathcal{R} be rich. Then, the minimum price Walrasian rules are the only rules on \mathcal{R}^n satisfying efficiency, strategy-proofness, individual rationality and no subsidy for losers.*

5 Main result

In this section, first we state the main theorem. Next, we explain how we prove the theorem.

5.1 Impossibility result

We consider rich domains containing some multi-demand preferences and we investigate whether *efficient* and *strategy-proof* rules still exist on such domains. In marked contrast to Fact 3 in Section 3, the results are negative. Namely, if there are more agents than objects, and if the domain is rich and contains even a single multi-demand preference relation, then no rule on the domain satisfies *efficiency*, *strategy-proofness*, *individual rationality* and *no subsidy for losers*.

Theorem *Let $n > m$. Let $R_0 \in \mathcal{R}^M$ and \mathcal{R} be a rich class of preferences such that $R_0 \in \mathcal{R}$. Then, no rule on \mathcal{R}^n satisfies efficiency, strategy-proofness, individual rationality and no subsidy for losers.*

Corollary 1 *Let $n > m$. Let $\mathcal{R} = \mathcal{R}^U \cup \mathcal{R}^M$. Then, no rule on \mathcal{R}^n satisfies efficiency, strategy-proofness, individual rationality and no subsidy for losers.*

Remark 4 The Corollary 1 is a standard form of impossibility results on *strategy-proofness* in that since Gibbard (1973) and Satterthwaite (1975), many impossibility results on *strategy-proofness* in this form are established. In such results, the domain is fixed and incompatibility of some properties of rules is established on this domain. The results of this form cannot be applied unless all the preferences in the fixed domain are deemed plausible. For example, the Corollary 1 cannot be applied unless all the preferences in \mathcal{R}^M in addition to \mathcal{R}^U are deemed plausible. On the other hand, our Theorem can be applied as soon as in addition to a rich domain \mathcal{R} , just one preference relation R_0 arbitrarily chosen from \mathcal{R}^M is deemed plausible. Accordingly our Theorem can be applied to more variety of environments than Corollary 1. For example, consider an environment where $n = 40$, $m = 20$ and there are only the preferences satisfying k -object-demand property for $k \in \{1, 2, 3, 4, 5\}$. The Theorem can be applied to this environment, but the Corollary 1 cannot be.

By Remark 3, we also have the following corollaries. These corollaries demonstrate the wide applicability of our results even more. In this paper, we do not maintain free disposal. However, it is a standard assumption for preferences. Corollary 2 states that our conclusion holds even if free disposal is assumed.

Corollary 2 *Let $n > m$. Let $R_0 \in \mathcal{R}^M$ and \mathcal{R} be such that $\mathcal{R} = \mathcal{R}_+^U \cup \{R_0\}$. Then, no rule on \mathcal{R}^n satisfies efficiency, strategy-proofness, individual rationality and no subsidy for losers.*

As Example 3 illustrates, free disposal is not a suitable assumption in some environment. Corollary 3 states that our conclusion holds even in such environment.

Corollary 3 *Let $n > m$. Let $R_0 \in \mathcal{R}^M$ and \mathcal{R} be such that $\mathcal{R} = (\mathcal{R}^U \setminus \mathcal{R}_+^U) \cup \{R_0\}$. Then, no rule on \mathcal{R}^n satisfies efficiency, strategy-proofness, individual rationality and no subsidy for losers.*

Remark 5 In this paper, we assume that preferences are drawn from a common class of \mathcal{R} . If the preferences of each agent are drawn from a class \mathcal{R}_i that depends on the identity of the agent, our theorem can be strengthened as follows: Suppose that, for each $i \in N$, \mathcal{R}_i is rich, and there are $j \in N$ and $R_j \in \mathcal{R}^M$ such that $R_j \in \mathcal{R}_j$. Then, when $n > m$, no rule on $\prod_{i \in N} \mathcal{R}_i$ satisfies *efficiency*, *strategy-proofness*, *individual rationality* and *no subsidy for losers*.

5.2 Sketch of the proof

5.2.1 Preliminary results

We state seven lemmas which we use in the sketch of the proof and in the formal proof. The proof of each lemma is relegated to the Appendix, or is omitted if it is straightforward.

Let $\mathcal{R} \subseteq \mathcal{R}^C$ be rich. Let f be a rule on \mathcal{R}^n satisfying *efficiency*, *strategy-proofness*, *individual rationality* and *no subsidy for losers*. Lemma 5 states that if an agent receives no object, then his payment is zero. This is immediate from *individual rationality* and *no subsidy for losers*. Thus we omit the proof.

Lemma 5 (Zero payment for losers) *Let $R \in \mathcal{R}^n$ and $i \in N$. If $A_i(R) = \emptyset$, $t_i(R) = 0$.*

Lemma 6 states that for each agent, his payment is at most the valuation, at $\mathbf{0}$, of the set of objects that he receives. This is immediate from *individual rationality*. Thus, we omit the proof.

Lemma 6 *For each $R \in \mathcal{R}^n$ and each $i \in N$, $t_i(R) \leq V_i(A_i(R); \mathbf{0})$.*

Lemma 7 states that each object is assigned to some agent. This follows from *efficiency*, $n > m$, and first object monotonicity. We omit the proof.

Lemma 7 (Full object assignment) *Let $n > m$. For each $R \in \mathcal{R}^n$ and each $a \in M$, there is $i \in N$ such that $a \in A_i(R)$.*

Lemma 8 is a necessary condition for *efficiency*.

Lemma 8 (Necessary condition for *efficiency*) *Let $R \in \mathcal{R}^n$ and $i, j \in N$ with $i \neq j$. Let $A_i, A_j \in \mathcal{M}$ be such that $A_i \cap A_j = \emptyset$ and $A_i \cup A_j \subseteq A_i(R) \cup A_j(R)$. Then, $V_i(A_i; f_i(R)) + V_j(A_j; f_j(R)) \leq t_i(R) + t_j(R)$.*

Although *no subsidy for losers* itself tells us nothing about payment levels for non-empty sets of objects, Lemma 9 states that for each non-empty set of object, there is a lower bound of the payment level for the set.

Lemma 9 (Payment lower bound) *Let $n > m$. Let $R \in \mathcal{R}^n$ and $i \in N$. Let $R'_i \in \mathcal{R} \cap \mathcal{R}^U$ be such that for each $a \in M$ and each $t_i \in \mathbb{R}$, $V'_i(a; (\emptyset, t_i)) - t_i < \min_{j \in N \setminus \{i\}} V_j(a; \mathbf{0})$.¹⁴ Then, $t_i(R) \geq V'_i(A_i(R); \mathbf{0})$.*

By first object monotonicity, Lemma 9 implies that for each $R \in \mathcal{R}^n$ and each $i \in N$, if $|A_i(R)| = 1$, then $t_i(R) \geq 0$.

Lemma 10 states that f coincides with an MPW rule on $(\mathcal{R} \cap \mathcal{R}^U)^n$. This is immediate from Fact 3 (ii). Thus we omit the proof.

Lemma 10 *Let $n > m$. For each $R \in (\mathcal{R} \cap \mathcal{R}^U)^n$, $f(R) \in Z_{\min}^W(R)$.*

¹⁴ Notice that in each class of preferences satisfying the richness, there exists such a preference relation.

Given $i \in N$ and $R_{-i} \in \mathcal{R}^{n-1}$, we define the **option set of agent i for R_{-i}** by

$$o_i(R_{-i}) \equiv \{z_i \in \mathcal{M} \times \mathbb{R} : \exists R_i \in \mathcal{R} \text{ s.t. } f_i(R_i, R_{-i}) = z_i\}.$$

Lemma 11 states that (i) the option set does not contain more than one bundle with the same set of objects, and (ii) each agent receives one of the most preferred bundles in his option set. This is straightforward from *strategy-proofness*. Thus, we omit the proof.

Lemma 11 *Let $i \in N$ and $R_{-i} \in \mathcal{R}^n$. (i) For each pair $(A_i, t_i), (A'_i, t'_i) \in o_i(R_{-i})$, if $A_i = A'_i$, then $t_i = t'_i$. (ii) For each $R_i \in \mathcal{R}$ and each $z_i \in o_i(R_{-i})$, $f_i(R_i, R_{-i}) R_i z_i$.*

5.2.2 Two-agent and three-object example

Since the proof of the Theorem is very complicated, we relegate it to the Appendix. Here we demonstrate the ideas and techniques of the proof by applying them to a particular example in a three-agent and two-object setting. Let $M = \{a, b\}$ and $N = \{1, 2, 3\}$. In the formal proof, the preference relation R_0 is an arbitrary element of \mathcal{R}^M , but here we pick R_0 from $\mathcal{R}^Q \cap \mathcal{R}^M$. For concreteness, let

$$v_0(a) = 20, v_0(b) = 18 \text{ and } v_0(\{a, b\}) = 40.$$

However, the idea of our proof does not depend on $R_0 \in \mathcal{R}^Q$. We assume $R_0 \in \mathcal{R}^Q$ only for simplicity of expression.

Let $\mathcal{R} \subseteq \mathcal{R}^C$ satisfy the richness and contain R_0 . For example, let $\mathcal{R} \supseteq \mathcal{R}^U \cup \{R_0\}$. We suppose that there is a rule f on \mathcal{R}^3 satisfying *efficiency*, *strategy-proofness*, *individual rationality* and *no subsidy for losers*, and derive a contradiction.

Step A: Constructing a preference profile.

Let $R_1 = R_0$. We construct $R_2 \in \mathcal{R}^U$ and $R_3 \in \mathcal{R}^U$ depending on R_1 so that a contradiction is derived. We define R_2 satisfying $V_2(a; \mathbf{0}) > v_1(\{a, b\})$ and

$$V_2(b; \mathbf{0}) - V_2(\emptyset; (b, 0)) < \min\{v_1(\{a, b\}) - v_1(a), v_1(\{a, b\}) - v_1(b)\}^{15}$$

¹⁵ In this sketch, we assume that $M = \{a, b\}$ and $R_0 \in \mathcal{R}^Q$ for the simplicity of expression. For a general multi-demand preference relation R_0 , the RHS of the first inequality is set as the maximal difference between various t_1 in $[0, V_1(\{a, b\}; \mathbf{0})]$ and the valuation of empty set at $(\{a, b\}, t_1)$ for R_1 , *i.e.*,

$$\max_{t_1 \in [0, V_1(\{a, b\}; \mathbf{0})]} \{t_1 - V_1(\emptyset; (\{a, b\}, t_1))\}.$$

For a general set M , the RHS is defined as \bar{t}_1 in the Appendix.

For a general multi-demand preference relation R_0 , the RHS of the second inequality is set as the minimum value of the marginal valuations of the second object, *i.e.*,

$$\min\left\{\min_{t_1 \in [0, V_1(\{a\}; \mathbf{0})]} V_1(\{a, b\}; (a, t_1)) - t_1, \min_{t_1 \in [0, V_1(\{b\}; \mathbf{0})]} V_1(\{a, b\}; (b, t_1)) - t_1\right\}.$$

For a general set M of objects, the RHS of the second inequality is defined as \underline{t}_1 in the Appendix.

For example, let $R_2 \in \mathcal{R}^U$ be such that for each $A_2 \in \mathcal{M} \setminus \{\emptyset\}$,

$$V_2(A_2; \mathbf{0}) = \begin{cases} 41 & \text{if } A_2 = \{a\}, \\ 12 & \text{if } A_2 = \{b\}, \\ 41 & \text{if } A_2 = \{a, b\}, \end{cases} \quad V_2(A_2; (\emptyset, -1)) = \begin{cases} 9 & \text{if } A_2 = \{a\}, \\ 10 & \text{if } A_2 = \{b\}, \\ 10 & \text{if } A_2 = \{a, b\}, \end{cases}$$

$$V_2(A_2, (\emptyset, -2)) = \begin{cases} -1 & \text{if } A_2 = \{a\}, \\ 9 & \text{if } A_2 = \{b\}, \\ 9 & \text{if } A_2 = \{a, b\}, \end{cases} \quad V_2(A_2; (\emptyset, -3)) = \begin{cases} -2 & \text{if } A_2 = \{a\}, \\ 0 & \text{if } A_2 = \{b\}, \\ 0 & \text{if } A_2 = \{a, b\}. \end{cases}$$

We define R_3 satisfying $R_3 \in \mathcal{R}^U \cap \mathcal{R}^Q$, and

$$v_3(a) = v_3(b) < \frac{3}{5} \min\{v_1(\{a, b\}) - v_1(a), v_1(\{a, b\}) - v_1(b)\}.$$

For example, let $v_3(a) = v_3(b) = 9$. Note that $V_2(a; \mathbf{0}) > v_3(\{a, b\})$. Let $R \equiv (R_1, R_2, R_3)$. Figure 4 illustrates R .

***** FIGURE 4 ($R = (R_1, R_2, R_3)$) ENTERS HERE *****

■

Step B: $A_2(R) \neq \emptyset$.

Suppose by contradiction that $A_2(R) = \emptyset$. By Lemma 5, $f_2(R) = \mathbf{0}$. By Lemma 7, there is $i \neq 2$ such that $a \in A_i(R)$.

Let $A_i = \emptyset$ and $A_2 = \{a\}$. Note that $A_i \cap A_2 = \emptyset$ and $A_i \cup A_2 \subseteq A_i(R) \cup A_2(R)$. If $i = 1$, then $V_1(\emptyset; f_1(R)) \geq t_1(R) - 40$. If $i = 3$, then $V_3(\emptyset; f_3(R)) = t_3(R) - 9$. Thus, $V_i(\emptyset; f_i(R)) \geq t_i(R) - 40$. Since $V_2(a; \mathbf{0}) = 41$,

$$\begin{aligned} V_i(A_i; f_i(R)) + V_2(A_2; f_2(R)) &= V_i(\emptyset; f_i(R)) + V_2(a; \mathbf{0}) && \text{(by } f_2(R) = \mathbf{0}) \\ &\geq t_i(R) - 40 + 41 \\ &> t_i(R) + t_2(R). && \text{(by } t_2(R) = 0) \end{aligned}$$

Thus, by Lemma 8, *efficiency* is violated, a contradiction. ■

Step C: $A_1(R) = a$.

Substep C-1: $(a, 9) \in o_1(R_{-1})$.

Let $R'_1 \in \mathcal{R}^U$ be such that

$$V'_1(a; \mathbf{0}) > \max\{V_2(a; \mathbf{0}), V_3(a; \mathbf{0})\} \text{ and } V'_1(b; \mathbf{0}) < \min\{V_2(b; \mathbf{0}), V_3(b; \mathbf{0})\}.$$

For example, let $R'_1 \in \mathcal{R}^U$ be such that

$$V'_1(a; \mathbf{0}) = 50, V'_1(b; \mathbf{0}) = 1 \text{ and } V'_1(\{a, b\}; \mathbf{0}) = 50.$$

Since $(R'_1, R_{-1}) \in (\mathcal{R}^U)^3$, by Lemma 10, $f(R'_1, R_{-1}) \in Z_{\min}^W(R'_1, R_{-1})$. Let $z \in Z$ be such that

$$z_1 = (a, 9), z_2 = (b, 9) \text{ and } z_3 = \mathbf{0}.$$

Figure 5 illustrates (R'_1, R_{-1}) and z .

***** FIGURE 5 ((R'_1, R_{-1}) and z in Step C) ENTERS HERE *****

Let $p \equiv (9, 9)$. Then, $D(R'_1, p) = \{\{a\}\}$, $D(R_2, p) = \{\{b\}\}$, and $D(R_3, p) = \{\emptyset, \{a\}, \{b\}\}$. Thus $(z, p) \in W(R'_1, R_{-1})$, implying $p_{\min}(R'_1, R_{-1}) \leq p$. If $p_{\min}^a(R'_1, R_{-1}) < 9$ or $p_{\min}^b(R'_1, R_{-1}) < 9$, then $0 \notin D(R'_1, p_{\min}(R'_1, R_{-1}))$ and for each $i \in \{2, 3\}$, $0 \notin D(R_i, p_{\min}(R'_1, R_{-1}))$, which implies $p_{\min}(R'_1, R_{-1}) \notin P(R'_1, R_{-1})$, a contradiction. Thus, $p_{\min}(R'_1, R_{-1}) = (9, 9)$. Moreover, z is the only WE allocation supported by $p_{\min}(R'_1, R_{-1})$. Thus, $f_1(R'_1, R_{-1}) = z$, and hence, $f_1(R'_1, R_{-1}) = (a, 9) \in o_1(R_{-1})$.

Substep C-2: $(b, 10) \in o_1(R_{-1})$.

Let $R''_1 \in \mathcal{R}^U$ be such that

$$V''_1(b; \mathbf{0}) > \max\{V_2(b; \mathbf{0}), V_3(b; \mathbf{0})\} \text{ and } V''_1(a; \mathbf{0}) < \min\{V_2(a; \mathbf{0}), V_3(a; \mathbf{0})\}.$$

For example, let $R''_1 \in \mathcal{R}^U$ be such that

$$V''_1(a; \mathbf{0}) = 1, V''_1(b; \mathbf{0}) = 50 \text{ and } V''_1(\{a, b\}; \mathbf{0}) = 50.$$

Since $(R''_1, R_{-1}) \in (\mathcal{R}^U)^3$, by Lemma 10, $f(R''_1, R_{-1}) \in Z_{\min}^W(R''_1, R_{-1})$. Let $z' \in Z$ be such that

$$z'_1 = (b, 10), z'_2 = (a, 9), \text{ and } z'_3 = \mathbf{0}.$$

Figure 6 illustrates (R''_1, R_{-1}) and z' .

***** FIGURE 6 ((R''_1, R_{-1}) and z' in Step C) ENTERS HERE *****

Let $p' \equiv (9, 10)$. Then, $D(R''_1, p') = \{\{b\}\}$, $D(R_2, p') = \{\{a\}, \{b\}\}$, and $D(R_3, p') = \{\emptyset, \{a\}\}$. Thus $(z', p') \in W(R''_1, R_{-1})$, implying $p_{\min}(R''_1, R_{-1}) \leq p'$. If $p_{\min}^a(R''_1, R_{-1}) < 9$, then $0 \notin D(R''_1, p_{\min}(R''_1, R_{-1}))$ and for each $i \in \{2, 3\}$, $0 \notin D(R_i, p_{\min}(R''_1, R_{-1}))$, which implies $p_{\min}(R''_1, R_{-1}) \notin P(R''_1, R_{-1})$, a contradiction. Thus, $p_{\min}^a(R''_1, R_{-1}) = 9$. If $p_{\min}^b(R''_1, R_{-1}) < 10$, then we have $D(R''_1, p_{\min}(R''_1, R_{-1})) = \{\{b\}\}$ and $D(R_2, p_{\min}(R''_1, R_{-1})) = \{\{b\}\}$, which further implies $p_{\min}(R''_1, R_{-1}) \notin P(R''_1, R_{-1})$, a contradiction. Thus, $p_{\min}(R''_1, R_{-1}) = (9, 10)$. Moreover, z' is the only WE allocation which is supported by $p_{\min}(R''_1, R_{-1})$. Thus, $f(R''_1, R_{-1}) = z'$, and hence, $f_1(R''_1, R_{-1}) = (b, 10) \in o_1(R_{-1})$.

Substep C-3: $A_1(R) = a$.

Since $A_2(R) \neq \emptyset$ by Step B, $|A_1(R)| \leq 1$. If $A_1(R) = \emptyset$, then by Lemma 5, we have $f_1(R) = \mathbf{0}$, and thus, $f_1(R'_1, R_{-1}) P_1 f_1(R)$, which contradicts *strategy-proofness*. Thus, $A_1(R) = a$ or b , and therefore, by (i) of Lemma 11, $f_1(R) = (a, 9)$ or $(b, 10)$. Since $(a, 9) P_1 (b, 10)$, (ii) of Lemma 11 implies $f_1(R) = (a, 9)$. ■

Step D: $f(R)$ is not efficient for R .

By $A_2(R) \neq \emptyset$ and $A_1(R) = a$, $A_2(R) = b$. By Lemma 6, $t_2(R) \leq V_2(b; \mathbf{0}) = 12$. By Lemma 9, $t_2(R) \geq 0$.

Let $z \equiv ((A_i, t_i))_{i \in N} \in Z$ be such that

$$z_1 = (\{a, b\}, 29), z_2 = (\emptyset, -3), \text{ and } z_3 = f_3(R).$$

Figure 7 illustrates z .

***** FIGURE 7 (z in Step D) ENTERS HERE *****

Since, $V_1(\{a, b\}; f_1(R)) = V_1(\{a, b\}; (a, 9)) = 29$, it is easy to see that $z_1 I_1 f_1(R)$ and $z_3 I_3 f_3(R)$. Also by $t_2(R) \geq 0$ and $A_2(R) = b$, $z_2 = (\emptyset, -3) I_2 (b, 0) R_2 f_2(R)$. Moreover, by $t_1(R) = 9$ and $t_2(R) \leq 12$,

$$\sum_{i \in N} t_i = 29 - 3 + t_3(R) = 26 + t_3(R) > \sum_{i \in M} t_i(R),$$

implying that $f(R)$ is not *efficient* for R , a contradiction. ■

We emphasize the difference between a (direct) proof of the Corollary 1 that one might write, and the proof of the Theorem that we have shown. To prove the Corollary 1 directly, we can freely pick preference profiles in $\mathcal{R}^U \cup \mathcal{R}^M$ to derive a contradiction. On the other hand, in the proof of the Theorem, we may only choose preferences from $\mathcal{R}^U \cup \{R_0\}$. Moreover, the preference relation R_0 , which could be anything in \mathcal{R}^M , forces us to construct profiles depending on R_0 , further complicating the process. In the above sketch, $R_0 \in \mathcal{R}^M$ is assumed to be quasi-linear, but the basic logic of the sketch works even in the case $R_0 \in \mathcal{R}^M \setminus \mathcal{R}^Q$.

In the formal proof in the Appendix, we have six steps. Steps A, B, C, and D correspond to Steps 1, 3, 4, and 6, respectively, in the formal proof. Steps 2 and 5 in the formal proof are necessary only for the more general case, so they do not appear in the above sketch.

6 Concluding remarks

In this article, we have considered an object assignment problem with money where each agent can receive more than one object. We focused on domains that contain enough variety of unit-demand preferences and some multi-demand preferences. We studied allocation rules satisfying *efficiency*, *strategy-proofness*, *individual rationality*, and *no subsidy for losers*, and showed that if the domain contains enough variety of unit-demand preferences and at least one multi-demand preference relation, and if there are more agents than objects, then no rule satisfies the four properties. As discussed in Section 1, we have been motivated by the search for *efficient* and *strategy-proof* rules on a domain which is not quasi-linear or unit-demand. Our result establishes the difficulty of designing *efficient* and *strategy-proof* rules on such a domain. We state three remarks on our result.

Maximal domain. Some literature on *strategy-proofness* investigates the existence of maximal domains on which there are rules satisfying desirable properties.¹⁶ A domain \mathcal{R}^n is a *maximal domain* for a list of properties of rules if there is a rule on \mathcal{R}^n satisfying the properties, and for each $\mathcal{R}' \supsetneq \mathcal{R}$, no rule on $(\mathcal{R}')^n$ satisfies the properties. Our result is rather closer to maximal domain results than impossibility results of the form of the Corollary 1. However, our result does not imply that the unit-demand domain is a maximal domain for the four properties in

¹⁶ For example, see Ching and Serizawa (1998), Berga and Serizawa (2000), Massó and Neme (2001), Ehlers (2002), etc.

the Theorem, since we add only multi-demand preferences to the rich domains and derive the non-existence of rules satisfying the four properties. In fact, what domains including $(\mathcal{R}^U)^n$ are maximal domains for the four properties is an open question.

However, we are sure that $(\mathcal{R}^U)^n$ is not a maximal domain for the four properties. For example, consider R_i in Example 4 and let $\mathcal{R} \equiv \mathcal{R}^U \cup \{R_i\}$. Since R_i does not satisfy the unit-demand property, $\mathcal{R} \not\supseteq \mathcal{R}^U$. Note that for each $R_j \in \mathcal{R}$, each $A_j \in \mathcal{M}$ with $|A_j| > 1$ and each $t_j \in \mathbb{R}$, if $t_j \geq 0$, then there is $a \in A_j$ such that $(a, t_j) I_j (A_j, t_j)$. Thus, since each agent never pays negative amount of money under the minimum price Walrasian rules, and since they satisfy the four properties on the unit-demand domain, they also satisfy the four properties on \mathcal{R}^n . Hence, $(\mathcal{R}^U)^n$ is not a maximal domain for the four properties.

Although we do not find maximal domains for the four properties, the multi-demand class includes most of natural preferences outside the unit-demand class. Thus, our result implies that on most of natural domains including the unit-demand domain, if there are more agents than objects, we have an impossibility of designing rules satisfying the four properties.

Other properties. *Efficiency* is not the only property studied in the literature on auction theory. For example, some authors study *strategy-proof* and *individually rational* rules that achieve as much revenue as possible. Since *efficiency* takes the auctioneer's revenue into account, *efficiency* is closely related to maximizing the auctioneer's revenue. However, there may exist *strategy-proof* and *individually rational* rules that is not *efficient* but achieve as much revenue as possible.¹⁷

While *efficiency* takes the auctioneer's revenue into account, some authors study another efficiency notion that takes only agents' preferences into account.¹⁸ An allocation is *efficient with no deficit* if (i) the sum of payments is nonnegative, and (ii) no other allocation with nonnegative sum of payments makes each agent at least as well off and at least one agent better off. Notice that *efficiency* is implied by *efficiency with no deficit*.¹⁹ Thus, the Theorem holds even if we replace *efficiency* by *efficiency with no deficit*.

Identical objects. Some literature on object assignment problems also study the case in which the objects are identical.²⁰ In this paper, we do not make this assumption. When objects are not identical, the domain includes a greater variety of preference profiles than when objects are identical. This variety plays an important role in our proof. Therefore, our theorem does not exclude the possibility that when objects are identical, multi-demand preferences can be added to the unit-demand domain without preventing the existence of rules satisfying the four properties.

¹⁷ For the single object case with quasi-linear preferences, Myerson type rules are not necessarily *efficient* but maximize the auctioneer's revenue. See Myerson (1981).

¹⁸ For example, see Sprumont (2013).

¹⁹ Let $R \equiv (R_i)_{i \in N}$ be a preference profile. Suppose there is an allocation $z \equiv ((A_i, t_i))_{i \in N}$ that is *efficiency with no deficit* for R but not *efficient* for R . Then, there is an allocation $z' \equiv ((A'_i, t'_i))_{i \in N}$ such that

$$\text{for each } i \in N, z'_i R_i z_i, \text{ for some } j \in N, z'_j P_j z_j, \text{ and } \sum_{i \in N} t'_i \geq \sum_{i \in N} t_i.$$

By condition (i) of *efficiency with no deficit*, we have $\sum_{i \in N} t_i \geq 0$. Thus, $\sum_{i \in N} t'_i \geq 0$. However, this implies that z is not *efficient with no deficit* for R , a contradiction.

²⁰ For example, see Saitoh and Serizawa (2008), Ashlagi and Serizawa (2012), Adachi (2014), etc.

Appendix: Proofs

A Proofs of Lemmas

Proof of Lemma 1: Since R_i is complete, transitive, and continuous, there is a continuous utility function $u_i : \mathcal{M} \times \mathbb{R} \rightarrow \mathbb{R}$ representing R_i . By possibility of compensation, there are $t'_i \in \mathbb{R}$ and $t''_i \in \mathbb{R}$ such that $(A'_i, t'_i) R_i z_i R_i (A'_i, t''_i)$, that is, $u_i(A'_i, t'_i) \geq u_i(z_i) \geq u_i(A'_i, t''_i)$. If we have either $(A'_i, t'_i) I_i z_i$ or $z_i I_i (A'_i, t''_i)$, then we are done. Thus, suppose $(A'_i, t'_i) P_i z_i$ and $z_i P_i (A'_i, t''_i)$, that is, $u_i(A'_i, t'_i) > u_i(z_i) > u_i(A'_i, t''_i)$.

Since u_i is continuous, $u_i(A'_i, \cdot)$ is continuous in \mathbb{R} . Moreover, by money monotonicity, $t'_i < t''_i$. Since $[t'_i, t''_i]$ is a closed interval, intermediate value theorem implies that there is $t_i^* \in (t'_i, t''_i)$ such that $u_i(A'_i, t_i^*) = u_i(z_i)$, that is $(A'_i, t_i^*) I_i z_i$. \square

Proof of (i) of Lemma 2: Suppose that there is $(A_i, t_i) \in D(R_i, p)$ such that $|A_i| > 1$. By $R_i \in \mathcal{R}^U$, there is $a \in A_i$ such that $(a, t_i) R_i (A_i, t_i)$. By $(A_i, t_i) \in B(p)$, $t_i = \sum_{b \in A_i} p^b$. By $p \in \mathbb{R}_{++}^m$ and $|A_i| > 1$, $p^a < \sum_{b \in A_i} p^b = t_i$. Thus, by money monotonicity,

$$(a, p^a) P_i (a, t_i) R_i (A_i, t_i),$$

which contradicts $(A_i, t_i) \in D(R_i, p)$. \square

Proof of (ii) of Lemma 2: Let $A'_i \in \mathcal{M}$. If $|A'_i| \leq 1$, then by the def. of A_i , $(A_i, \sum_{a \in A_i} p^a) R_i (A'_i, \sum_{a \in A'_i} p^a)$. Suppose $|A'_i| > 1$. By $R_i \in \mathcal{R}^U$, there is $a \in A'_i$ such that $(a, p^a) R_i (A_i, \sum_{b \in A'_i} p^b)$. By $p^a \geq 0$ and by money monotonicity, $(a, p^a) R_i (a, \sum_{b \in A'_i} p^b) R_i (A'_i, \sum_{b \in A'_i} p^b)$. Thus, by the def. of A_i ,

$$(A_i, \sum_{b \in A_i} p^b) R_i (a, p^a) R_i (A'_i, \sum_{b \in A'_i} p^b).$$

Thus, $(A_i, \sum_{b \in A_i} p^b) \in D(R_i, p)$. \square

Proof of (i) of Lemma 3: By contradiction, suppose that $n > m$ and $p^a = 0$ for some $a \in M$. Then, by money monotonicity, for each $i \in N$, $(a, p^a) P_i \mathbf{0}$. Thus, for each $i \in N$, $\mathbf{0} \notin D(R_i, p)$, which implies that for each $((A, t), p) \in W(R)$, $A_i \neq \emptyset$. However, this contradicts $n > m$. \square

Proof of (ii) of Lemma 3: By $p \in P(R)$, there is $(A, t) \in Z$ such that $((A, t), p) \in W(R)$.

Let $N^* = \{i \in N : |A_i| > 1\}$ and $i \in N^*$. By $R_i \in \mathcal{R}^U$, there is $a \in A_i$ such that $(a, t_i) R_i (A_i, t_i)$. By $(A_i, t_i) \in B(p)$, $t_i = \sum_{b \in A_i} p^b$. By $p \in \mathbb{R}_+^m$, $p^a \leq \sum_{b \in A_i} p^b = t_i$. Thus, by money monotonicity, $(a, p^a) R_i (a, t_i) R_i (A_i, t_i)$, which implies $(a, p^a) \in D(R_i, p)$. Hence, for each $i \in N^*$, there is $a_i \in A_i$ such that $(a_i, p^{a_i}) \in D(R_i, p)$.

Let $z' \in Z$ be such that for each $i \in N^*$, $z'_i = (a_i, p^{a_i})$, and for each $i \in N \setminus N^*$, $z'_i = (A_i, t_i)$. Then, each agent receives at most one object at z' , and clearly, $(z', p) \in W(R)$. \square

Proof of Lemma 4: Suppose by contradiction that $R_i \in \mathcal{R}^U$ and $|A_i(R)| > 1$. Then, there is $a \in A_i(R)$ such that $(a, t_i(R)) R_i f_i(R)$. By $|A_i(R)| > 1$, there is $b \in A_i(R)$ such that $b \neq a$.

By $n > m$, there is $j \in N \setminus \{i\}$ such that $A_j(R) = \emptyset$. Let $z \equiv ((A_k, t_k))_{k \in N} \in Z$ be such that

$$\begin{aligned} z_i &= (a, t_i(R)), \\ z_j &= (b, t_j(R)), \text{ and} \\ z_k &= f_k(R) \text{ for each } k \in N \setminus \{i, j\}. \end{aligned}$$

Clearly, $\sum_{k \in N} t_k = \sum_{k \in N} t_k(R)$, and for each $k \in \setminus \{i, j\}$, $z_k \succ_k f_k(R)$. Moreover, $z_i = (a, t_i(R)) \succ_{R_i} f_i(R)$, and by first object monotonicity, $z_j = (b, t_j(R)) \succ_{P_j} f_j(R)$. This contradicts *efficiency*. \square

Proof of Lemma 8: Suppose by contradiction that $V_i(A_i; f_i(R)) + V_j(A_j; f_j(R)) > t_i(R) + t_j(R)$. Let $z' \in Z$ be such that $z'_i = (A_i, V_i(A_i; f_i(R)))$, $z'_j = (A_j, V_j(A_j; f_j(R)))$, and for each $k \in N \setminus \{i, j\}$, $z'_k = f_k(R)$. Then $z'_k \succ_k f_k(R)$ for each $k \in N$. Moreover, $V_i(A_i; f_i(R)) + V_j(A_j; f_j(R)) + \sum_{k \neq i, j} t_k(R) > \sum_{k \in N} t_k(R)$. By Remark 2, this contradicts *efficiency*. \square

Proof of Lemma 9: (Figure 8.) Suppose by contradiction that $t_i(R) < V'_i(A_i(R); \mathbf{0})$. If $A_i(R) = \emptyset$, then $t_i(R) < V'_i(\emptyset; \mathbf{0}) = 0$, which contradicts *no subsidy for losers*. Hence, $A_i(R) \neq \emptyset$.

Next, we show $A_i(R'_i, R_{-i}) \neq \emptyset$. Suppose not. Then, by Lemma 5, $f_i(R'_i, R_{-i}) = \mathbf{0}$. By $t_i(R) < V'_i(A_i(R); \mathbf{0})$, $f_i(R) \succ_{P'_i} \mathbf{0} = f_i(R'_i, R_{-i})$, which contradicts *strategy-proofness*. Hence $A_i(R'_i, R_{-i}) \neq \emptyset$.

By $R'_i \in \mathcal{R}^U$, $A_i(R'_i, R_{-i}) \neq \emptyset$, and Lemma 4, there is $a \in M$ such that $A_i(R'_i, R_{-i}) = a$. Since $n > m$ and $A_i(R'_i, R_{-i}) \neq \emptyset$, there is $j \in N \setminus \{i\}$ such that $A_j(R'_i, R_{-i}) = \emptyset$. By Lemma 5, $f_j(R'_i, R_{-i}) = \mathbf{0}$. Thus, letting $s_i \equiv V'_i(\emptyset; f_i(R'_i, R_{-i}))$,

$$\begin{aligned} &V'_i(\emptyset; f_i(R'_i, R_{-i})) + V_j(a; f_j(R'_i, R_{-i})) \\ &= s_i + V_j(a; \mathbf{0}) && \text{(by } s_i = V'_i(\emptyset; f_i(R'_i, R_{-i})) \text{ and } f_j(R'_i, R_{-i}) = \mathbf{0}) \\ &= t_i(R'_i, R_{-i}) - (t_i(R'_i, R_{-i}) - s_i) + V_j(a; \mathbf{0}) \\ &= t_i(R'_i, R_{-i}) - (V'_i(a; (\emptyset, s_i)) - s_i) + V_j(a; \mathbf{0}) && \text{(by } V'_i(a; (\emptyset, s_i)) = t_i(R'_i, R_{-i})) \\ &> t_i(R'_i, R_{-i}) && \text{(by } V'_i(a; (\emptyset, s_i)) - s_i < V_j(a; \mathbf{0}) \\ &= t_i(R'_i, R_{-i}) + t_j(R'_i, R_{-i}). && \text{(by } t_j(R'_i, R_{-i}) = 0) \end{aligned}$$

This contradicts Lemma 8. \square

***** FIGURE 8 (Illustration of proof of Lemma 9) ENTERS HERE *****

B Proof of Theorem

The proof of the Theorem has six steps.

Step 1: Constructing preferences.

Let $R_1 \equiv R_0$. For each $a \in M$, let $\mathcal{M}_a \equiv \{A_1 \in \mathcal{M} : a \in A_1\}$.

Claim 1: There is $\underline{t}_1 \in \mathbb{R}$ such that $\underline{t}_1 > 0$ and

$$\underline{t}_1 = \min_{a \in M} \min_{t_1 \in [0, V_1(\{a\}; \mathbf{0})]} \max_{A_1 \in \mathcal{M}_a} \{V_1(A_1; (a, t_1)) - t_1\}.$$

Proof: For each $a \in M$, let g^a be a function on $\mathcal{M}_a \times \mathbb{R}$ such that for each $A_1 \in \mathcal{M}_a$ and each $t_1 \in \mathbb{R}$, $g^a(A_1, t_1) = V_1(A_1; (a, t_1)) - t_1$, and let g_{\max}^a be a function on \mathbb{R} such that for each $t_1 \in \mathbb{R}$, $g_{\max}^a(t_1) = \max_{A_1 \in \mathcal{M}_a} g^a(A_1, t_1)$.

Note that for each $a \in M$, and each $A_1 \in \mathcal{M}_a$, $g^a(A_1, \cdot)$ is continuous in \mathbb{R} . Thus, Berge's maximum theorem implies that for each $a \in M$, $g_{\max}^a(\cdot)$ is also continuous in \mathbb{R} .²¹ For each $a \in M$, since $[0, V_1(a; \mathbf{0})]$ is compact, there is $\hat{t}_1^a \in [0, V_1(a; \mathbf{0})]$ such that $g_{\max}^a(\hat{t}_1^a) = \min_{t_1 \in [0, V_1(a; \mathbf{0})]} g_{\max}^a(t_1)$. For each $a \in M$, let $\underline{t}_1^a \equiv g_{\max}^a(\hat{t}_1^a)$. Since M is finite, $\min\{\underline{t}_1^a : a \in M\}$ exists. Let $\underline{t}_1 \equiv \min\{\underline{t}_1^a : a \in M\}$. Then,

$$\underline{t}_1 = \min_{a \in M} \min_{t_1 \in [0, V_1(a; \mathbf{0})]} \max_{A_1 \in \mathcal{M}_a} \{V_1(A_1; (a, t_1)) - t_1\}.$$

Next, we show $\bar{t}_1 > 0$. Let $a \in M$ be such that $\underline{t}_1 = \max_{A_1 \in \mathcal{M}_a} \{V_1(A_1; (a, \hat{t}_1^a)) - \hat{t}_1^a\}$. By $R_1 \in \mathcal{R}^M$, there is $\hat{A}_1 \in \mathcal{M}_a$ such that $(\hat{A}_1, \hat{t}_1^a) P_1(a, \hat{t}_1^a)$, that is, $V_1(\hat{A}_1; (a, \hat{t}_1^a)) - \hat{t}_1^a > 0$. Thus,

$$\underline{t}_1 = \max_{A_1 \in \mathcal{M}_a} \{V_1(A_1; (a, \hat{t}_1^a)) - \hat{t}_1^a\} \geq V_1(\hat{A}_1; (a, \hat{t}_1^a)) - \hat{t}_1^a > 0.$$

□

By Claim 1, there is $d_* \in \mathbb{R}_{++}$ such that $5(m-1)d_* < \underline{t}_1$ and for each $a \in M$, $(a, 3d_*) P_1 \mathbf{0}$. Let $a^* \in M$ be such that for each $a \in M \setminus \{a^*\}$, $(a^*, 3d_*) R_1(a, 3d_*)$. Without loss of generality, assume $a^* = 1$.

Since \mathcal{R} is rich, there is $R'_1 \in \mathcal{R} \cap \mathcal{R}^U$ such that for each $a \in M$ and each $t_1 \in \mathbb{R}$, $V'_1(a; (\emptyset, t_1)) = \frac{d_*}{2} + t_1$. Let

$$s_1^* \equiv \min\{0, \min_{A_1 \in \mathcal{M} \setminus \{\emptyset\}} V'_1(A_1; \mathbf{0})\}, \text{ and}$$

$$\mathcal{M}(R_1, s_1^*) \equiv \{A_1 \in \mathcal{M} \setminus \{\emptyset\} : V_1(A_1; \mathbf{0}) \geq s_1^*\}.$$

Note that by first object monotonicity, $\mathcal{M}(R_1, s_1^*) \neq \emptyset$.

Claim 2: There is $\bar{t}_1 \in \mathbb{R}$ such that $\bar{t}_1 \geq \underline{t}_1$ and

$$\bar{t}_1 = \max_{A_1 \in \mathcal{M}(R_1, s_1^*)} \max_{t_1 \in [s_1^*, V_1(A_1; \mathbf{0})]} \{t_1 - V_1(\emptyset; (A_1, t_1))\}.$$

Proof: For each $A_1 \in \mathcal{M}(R_1, s_1^*)$, let g^{A_1} be a function on \mathbb{R} such that for each $t_1 \in \mathbb{R}$, $g^{A_1}(t_1) = t_1 - V_1(\emptyset; (A_1, t_1))$. Note that for each $A_1 \in \mathcal{M}(R_1, s_1^*)$, g^{A_1} is continuous in \mathbb{R} . For each $A_1 \in \mathcal{M}(R_1, s_1^*)$, since $[s_1^*, V_1(A_1; \mathbf{0})]$ is compact, there is $\hat{t}_1^{A_1} \in [s_1^*, V_1(A_1; \mathbf{0})]$ such that $g^{A_1}(\hat{t}_1^{A_1}) = \max_{t_1 \in [s_1^*, V_1(A_1; \mathbf{0})]} g^{A_1}(t_1)$. For each $A_1 \in \mathcal{M}(R_1, s_1^*)$, let $\bar{t}_1^{A_1} = g^{A_1}(\hat{t}_1^{A_1})$. Since $\mathcal{M}(R_1, s_1^*)$ is finite, $\max\{\bar{t}_1^{A_1} : A_1 \in \mathcal{M}(R_1, s_1^*)\}$ exists. Let $\bar{t}_1 \equiv \max\{\bar{t}_1^{A_1} : A_1 \in \mathcal{M}(R_1, s_1^*)\}$.

²¹ See Berge (1963).

Then,

$$\bar{t}_1 = \max_{A_1 \in \mathcal{M}(R_1, s_1^*)} \max_{t_1 \in [s_1^*, V_1(A_1; \mathbf{0})]} \{t_1 - V_1(\emptyset; (A_1, t_1))\}.$$

Next, we show $\bar{t}_1 \geq \underline{t}_1$. Let $a \in M$, $t_1 \in [0, V_1(a; \mathbf{0})]$ and $A_1 \in \mathcal{M}_a$ be such that $\underline{t}_1 = V_1(A_1; (a, t_1)) - t_1$. Let $\hat{t}_1 \equiv V_1(A_1; (a, t_1))$. Since $V_1(\emptyset; (A_1, \hat{t}_1)) = V_1(\emptyset; (a, t_1))$ and since first object monotonicity implies $t_1 - V_1(\emptyset; (a, t_1)) > 0$,

$$\begin{aligned} \hat{t}_1 - V_1(\emptyset; (A_1, \hat{t}_1)) &= V_1(A_1; (a, t_1)) - t_1 + t_1 - V_1(\emptyset; (a, t_1)) \\ &> V_1(A_1; (a, t_1)) - t_1. \end{aligned}$$

By $t_1 \leq V_1(a; \mathbf{0})$, $\hat{t}_1 \leq V_1(A_1; \mathbf{0})$. By $t_1 \geq 0$, $\underline{t}_1 > 0$, and $s_1^* \leq 0$, $\hat{t}_1 = \underline{t}_1 + t_1 > 0 \geq s_1^*$. Thus, $s_1^* \leq \hat{t}_1 \leq V_1(A_1; \mathbf{0})$. This implies $A_1 \in \mathcal{M}(R_1, s_1^*)$, and thus,

$$\begin{aligned} \bar{t}_1 &= \max_{A'_1 \in \mathcal{M} \setminus \{\emptyset\}} \max_{t'_1 \in [0, V_1(A'_1; \mathbf{0})]} \{t'_1 - V_1(\emptyset; (A'_1, t'_1))\} \\ &\geq \hat{t}_1 - V_1(\emptyset; (A_1, \hat{t}_1)) \\ &\geq V_1(A_1; (a, t_1)) - t_1 = \underline{t}_1. \end{aligned}$$

□

Let $d^* \in \mathbb{R}_{++}$ be such that $d^* > \bar{t}_1$. Note that by Claim 2, $5(m-1)d_* < \underline{t}_1 \leq \bar{t}_1 < d^*$. Since \mathcal{R} is rich, for each $i \in \{2, \dots, m\}$, there is $R_i \in \mathcal{R} \cap \mathcal{R}^U$ satisfying the following conditions:

$$V_i(a; \mathbf{0}) = \begin{cases} d^* & \text{if } a = i-1, \\ 4d_* & \text{if } a = i, \\ d_* & \text{otherwise,} \end{cases} \quad (ia)$$

$$V_i(\emptyset; (i, 0)) = -d_*, \text{ and} \quad (ib)$$

$$\text{for each } a \in M \setminus \{i\}, V_i(a; (i, 3d_*)) < 0. \quad (ic)$$

Since \mathcal{R} is rich, for each $i \in N \setminus \{1, \dots, m\}$, there is $R_i \in \mathcal{R} \cap \mathcal{R}^U$ such that for each $a \in M$ and each $t_i \in \mathbb{R}$,

$$V_i(a; (\emptyset, t_i)) = 3d_* + t_i.$$

Denote $R \equiv (R_1, \dots, R_n)$. Figure 9 illustrates R_i and R_j , where $i \in \{2, \dots, m\}$ and $j \in \{m+1, \dots, n\}$.

***** FIGURE 9 (Illustrations of R_i ($i \in \{2, \dots, m\}$) and R_j ($j \in \{m+1, \dots, m\}$)).
ENTERS HERE *****

Notice that for each $a \in M$ and each $t_1 \in \mathbb{R}$,

$$V_1'(a; (\emptyset, t_1)) - t_1 = \frac{d_*}{2} < \min_{i \in N \setminus \{1\}} V_i(a; \mathbf{0}).$$

Thus, by Lemma 9, $t_1(R) \geq V_1'(A_1(R); \mathbf{0}) \geq s_1^*$. ■

Step 2: For each $i \in \{2, \dots, m\}$, $V_i(\emptyset; f_i(R)) \geq -d_$.*

Let $i \in \{2, \dots, m\}$. If $A_i(R) = \emptyset$, then by Lemma 5, $V_i(\emptyset; f_i(R)) = t_i(R) = 0 > -d_*$. Suppose $A_i(R) \neq \emptyset$. By $R_i \in \mathcal{R}^U$ and Lemma 4, there is $a \in M$ such that $A_i(R) = a$. Thus, Lemma 9 implies $t_i(R) \geq 0$.

If $a = i$, then by money monotonicity, $(i, 0) R_j f_j(R)$, and thus, by ib , $V_i(\emptyset; f_i(R)) \geq V_i(\emptyset; (i, 0)) = -d_*$. If $a \neq i$, then by ic ,

$$V_i(a; (i, 0)) < V_i(a; (i, 3d_*)) < 0 \leq t_i(R),$$

which implies $(i, 0) P_i f_i(R)$, and thus, by ib $V_i(\emptyset; f_i(R)) > V_i(\emptyset; (i, 0)) = -d_*$. ■

Step 3: For each $i \in \{2, \dots, m\}$, $A_i(R) \neq \emptyset$.

Suppose by contradiction that there is an agent $i \in \{2, \dots, m\}$ such that $A_i(R) = \emptyset$. By Lemma 5, $t_i(R) = 0$. By Lemma 7, there is an agent $j \in N \setminus \{i\}$ such that $i - 1 \in A_j(R)$. We show the following claim.

Claim 1: $t_j(R) - V_j(\emptyset; f_j(R)) < d^$.*

Proof: We have three cases.

Case 1: $j = 1$. By Lemma 6 and Lemma 9, $s_1^* \leq t_1(R) \leq V_1(A_1(R); \mathbf{0})$, implying $A_1(R) \in \mathcal{M}(R_1, s_1^*)$. Thus, by the def. of \bar{t}_1 ,

$$t_1(R) - V_1(\emptyset; f_1(R)) \leq \bar{t}_1 < d^*.$$

Case 2: $j \in \{2, \dots, m\}$. By $R_j \in \mathcal{R}^U$ and Lemma 4, $A_j(R) = i - 1$. By Lemma 6, $t_j(R) \leq V_j(i - 1; \mathbf{0})$. Since $j - 1 \neq i - 1$, $V_j(i - 1; \mathbf{0}) = 4d_*$ or d_* . Thus, $V_j(i - 1; \mathbf{0}) \leq 4d_*$. Moreover, by Step 2, $V_j(\emptyset; f_j(R)) \geq -d_*$. Therefore, by $5d_* < 5(m - 1)d_* < d^*$,

$$t_j(R) - V_j(\emptyset; f_j(R)) \leq V_j(i - 1; \mathbf{0}) + d_* \leq 5d_* < d^*.$$

Case 3: $j \in \{m + 1, \dots, n\}$. By $R_j \in \mathcal{R}^U$ and Lemma 4, $A_j(R) = i - 1$. Let $t_j^* \equiv V_j(\emptyset; f_j(R))$. By the def. of R_j and $3d_* < 5(m - 1)d_* < d^*$,

$$t_j(R) - V_j(\emptyset; f_j(R)) = V_j(i - 1; (\emptyset, t_j^*)) - t_j^* = 3d_* < d^*.$$

□

By $f_i(R) = \mathbf{0}$, $V_i(i - 1; f_i(R)) = V_i(i - 1; \mathbf{0}) = d^*$. Thus, by Claim 1,

$$V_i(i - 1; f_i(R)) + V_j(\emptyset; f_j(R)) > d^* + t_j(R) - d^* = t_i(R) + t_j(R).$$

This contradicts Lemma 8. ■

Step 4: $A_1(R) = 1$.

First we show the following claim.

Claim 1: For each $a \in M$, there is $t_1(a) \in \mathbb{R}$ such that $(a, t_1(a)) \in o_1(R_{-1})$ and

$$t(a) \begin{cases} \leq 3d_* & \text{if } a = 1, \\ > 3d_* & \text{otherwise.} \end{cases}$$

Proof: Let $a \in M$. Since \mathcal{R} is rich, there is $R_1'' \in \mathcal{R} \cap \mathcal{R}^U$ such that

$$V_1''(a; \mathbf{0}) > \max_{i \in N} V_i(a; \mathbf{0}), \quad (1a'')$$

$$\text{for each } b \in M \setminus \{a\}, V_1''(b; \mathbf{0}) < \min_{i \in N} V_i(b; \mathbf{0}), \text{ and} \quad (1b'')$$

$$\text{for each } b \in M \setminus \{a\}, V_1''(b; (a, 3d_*)) < 0. \quad (1c'')$$

Figure 10 illustrates R_1'' .

***** FIGURE 10 (Illustration of R_1'') ENTERS HERE *****

By $(R_1'', R_{-1}) \in (\mathcal{R}^U)^n$ and Lemma 10, $f(R_1'', R_{-1}) \in Z_{\min}^W(R_1'', R_{-1})$. Let $p \equiv p_{\min}(R_1'', R_{-1})$. By $(R_1'', R_{-1}) \in (\mathcal{R}^U)^n$ and Lemma 4, for each $i \in N$, $|A_i(R_1'', R_{-1})| \leq 1$. In the following three paragraphs, we show $(a, p^a) \in o_1(R_{-1})$.

First, suppose $A_1(R_1'', R_{-1}) = \emptyset$. By Lemma 5, $f_1(R_1'', R_{-1}) = \mathbf{0}$. By $A_1(R_1'', R_{-1}) = \emptyset$ and Lemma 7, there is an agent $i \in N \setminus \{1\}$ such that $A_i(R_1'', R_{-1}) = a$. Since $f(R_1'', R_{-1}) \in Z_{\min}^W(R_1'', R_{-1})$, $t_i(R_1'', R_{-1}) = p^a$. By Lemma 6, $p^a = t_i(R_1'', R_{-1}) \leq V_i(a; \mathbf{0})$. Since $f(R_1'', R_{-1}) \in Z_{\min}^W(R_1'', R_{-1})$, $f_1(R_1'', R_{-1}) = \mathbf{0} \in D(R_1'', p)$, and thus, $\mathbf{0} R_1''(a, p^a)$. Therefore,

$$V_1''(a; \mathbf{0}) \leq p^a \leq V_i(a; \mathbf{0}).$$

This contradicts $1a''$. Hence, $A_1(R_1'', R_{-1}) \neq \emptyset$.

Next, suppose that for some $b \in M \setminus \{a\}$, $A_1(R_1'', R_{-1}) = b$. Since $f(R_1'', R_{-1}) \in Z_{\min}^W(R_1'', R_{-1})$, $t_1(R_1'', R_{-1}) = p^b$. By Lemma 6, $p^b = t_1(R_1'', R_{-1}) \leq V_1''(b; \mathbf{0})$. By $A_1(R_1'', R_{-1}) \neq \emptyset$ and $n > m$, there is an agent $i \in N \setminus \{1\}$ such that $A_i(R_1'', R_{-1}) = \emptyset$. By Lemma 5, $f_i(R_1'', R_{-1}) = \mathbf{0}$. Since $f(R_1'', R_{-1}) \in Z_{\min}^W(R_1'', R_{-1})$, $f_i(R_1'', R_{-1}) = \mathbf{0} \in D(R_i, p)$, and thus, $\mathbf{0} R_i(b, p^b)$. Therefore,

$$V_i(b; \mathbf{0}) \leq p^b \leq V_1''(b; \mathbf{0}).$$

This contradicts $1b''$. Thus for each $b \in M \setminus \{a\}$, $A_1(R_1'', R_{-1}) \neq b$.

By $A_1(R_1'', R_{-1}) \neq \emptyset$, $A_1(R_1'', R_{-1}) \neq b$ for each $b \in M \setminus \{a\}$, and $|A_1(R_1'', R_{-1})| \leq 1$, we conclude that $A_1(R_1'', R_{-1}) = a$. Since $f(R_1'', R_{-1}) \in Z_{\min}^W(R_1'', R_{-1})$, $t_1(R_1'', R_{-1}) = p^a$. Hence, $(a, p^a) \in o_1(R_{-1})$.

Next, we show that $p^a \leq 3d_*$ if $a = 1$, and $p^a > 3d_*$ otherwise.

Case 1: $a = 1$. Let $z \in Z$ be such that for each $i \in N$,

$$z_i = \begin{cases} (i, 3d_*) & \text{if } i \in \{1, \dots, m\} \\ \mathbf{0} & \text{otherwise.} \end{cases}$$

Let $\hat{p} \in \mathbb{R}_+^m$ be such that for each $b \in M$, $\hat{p}^b = 3d_*$. We show that $z_1 \in D(R_1'', \hat{p})$, and for each $i \in N \setminus \{1\}$, $z_i \in D(R_i, \hat{p})$. This implies that $(z, \hat{p}) \in W(R_1'', R_{-1})$, and thus, by

$p = p_{\min}(R''_1, R_{-1})$, we conclude that $p^1 \leq \hat{p}^1 = 3d_*$.

Note that by $\hat{p} \in \mathbb{R}_{++}^m$ and (i) of Lemma 2, for each $(A_1, t_1) \in D(R''_1, \hat{p})$, $|A_1| \leq 1$, and for each $i \in N \setminus \{1\}$ and each $(A_i, t_i) \in D(R_i, \hat{p})$, $|A_i| \leq 1$.

Let $i \in N$. We have three subcases.

Subcase 1-1: $i = 1$. By $1c''$ and $\hat{p} \in \mathbb{R}_{++}^m$, for each $b \in M \setminus \{1\}$,

$$V_1''(b; (1, \hat{p}^1)) = V_1''(b; (1, 3d_*)) < 0 < \hat{p}^b.$$

Thus for each $b \in M \setminus \{1\}$, $(1, \hat{p}^1) P_1'(b, \hat{p}^b)$. Also by $1c''$ and first object monotonicity, $V_1''(\emptyset; (1, \hat{p}^1)) = V_1''(\emptyset; (1, 3d_*)) < 0$, which implies $(1, \hat{p}^1) P_1' \mathbf{0}$. Thus, $z_1 = (1, \hat{p}^1) \in D(R''_1, \hat{p})$.

Subcase 1-2: $i \in \{2, \dots, m\}$. By ic and $\hat{p} \in \mathbb{R}_{++}^m$, for each $b \in M \setminus \{i\}$,

$$V_i(b; (i, \hat{p}^i)) = V_i(b; (i, 3d_*)) < 0 < \hat{p}^b.$$

Thus, for each $b \in M \setminus \{i\}$, $(i, \hat{p}^i) P_i(b, \hat{p}^b)$. Also by ic and first object monotonicity, $V_i(\emptyset; (i, \hat{p}^i)) = V_i(\emptyset; (i, 3d_*)) < 0$, which implies $(i, \hat{p}^i) P_i \mathbf{0}$. Thus, $z_i = (i, \hat{p}^i) \in D(R_i, \hat{p})$.

Subcase 1-3: $i \in \{m+1, \dots, n\}$. For each $b \in M$, $V_i(b; \mathbf{0}) = 3d_* = \hat{p}^b$. This implies $\mathbf{0} R_i(b, \hat{p}^b)$. Thus, $z_i = \mathbf{0} \in D(R_i, \hat{p})$.

Case 2: $a \in \{2, \dots, m\}$. Let $i = a$. Suppose by contradiction that $p^a \leq 3d_*$. Note that by $n < m$ and (i) of Lemma 3, $p \in \mathbb{R}_{++}^m$. Thus by (i) of Lemma 2, for each $(A_i, t_i) \in D(R_i, p)$, $|A_i| \leq 1$. By $p^a \leq 3d_*$, $i = a$, ic , and $p \in \mathbb{R}_{++}^m$, for each $b \in M \setminus \{a\}$,

$$V_i(b; (a, p^a)) \leq V_i(b; (i, 3d_*)) < 0 < p^b.$$

Thus for each $b \in M \setminus \{a\}$, $(a, p^a) P_i(b, p^b)$. Also by $i = a$, $p^a \leq 3d_*$, ic , and first object monotonicity, $V_i(\emptyset; (a, p^a)) \leq V_i(\emptyset; (i, 3d_*)) < 0$, which implies $(a, p^a) P_i \mathbf{0}$. Therefore, $D(R_i, p) = \{(a, p^a)\}$. Since $f(R''_1, R_{-1}) \in Z_{\min}^W(R''_1, R_{-1})$, $A_i(R''_1, R_{-1}) = a$. This contradicts $A_1(R''_1, R_{-1}) = a$. \square

Recall that $(1, 3d_*) P_1 \mathbf{0}$ and for each $a \in M \setminus \{1\}$, $(1, 3d_*) R_1(a, 3d_*)$. By Claim 1 of Step 4, $t(1) \leq 3d_*$ and for each $a \in M \setminus \{1\}$, $t(a) > 3d_*$. Thus,

$$(1, t(1)) R_1(1, 3d_*) P_1 \mathbf{0},$$

and for each $a \in M \setminus \{1\}$,

$$(1, t(1)) R_1(1, 3d_*) R_1(a, 3d_*) P_1(a, t(a)).$$

Therefore, (ii) of Lemma 11 implies $A_1(R) \neq \emptyset$ and for each $a \in M \setminus \{1\}$, $A_1(R) \neq a$. By Step 3, $|A_1(R)| \leq 1$, because otherwise there exists an agent $i \in \{2, \dots, m\}$ such that $A_i(R) = \emptyset$, which contradicts Step 3. Hence, $A_1(R) = 1$. \blacksquare

Step 5: For each $i \in \{2, \dots, m\}$, $A_i(R) = i$.

We show that for each $i \in \{2, \dots, m\}$, $A_i(R) \subseteq \{i, i-1\}$. Then, since $A_1(R) = 1$ by Step 4, we conclude that for each $i \in \{2, \dots, m\}$, $A_i(R) = i$.

Suppose by contradiction that there is $i \in \{2, \dots, m\}$ such that $A_i(R) \not\subseteq \{i-1, i\}$. By Step 3 and Lemma 4, there is $a \in M \setminus \{i-1, i\}$ such that $A_i(R) = a$. By $a \in M \setminus \{i-1, i\}$ and ia , $V_i(a; \mathbf{0}) = d_*$. Thus, by Step 2 and Lemma 6,

$$V_i(\emptyset; f_i(R)) \geq -d_* = V_i(a; \mathbf{0}) - 2d_* \geq t_i(R) - 2d_*. \quad (1)$$

Let $j \in \{m+1, \dots, n\}$. By Step 3 and Step 4, $A_j(R) = \emptyset$, and thus, by Lemma 5, $f_j(R) = \mathbf{0}$. Thus, by the def. of R_j ,

$$V_j(a; f_j(R)) = V_j(a; \mathbf{0}) = 3d_*. \quad (2)$$

Therefore, by (1), (2), and $t_j(R) = 0$,

$$V_i(\emptyset; f_i(R)) + V_j(a; f_j(R)) \geq t_i(R) - 2d_* + 3d_* > t_i(R) = t_i(R) + t_j(R).$$

This contradicts Lemma 8. ■

Step 6: Completing the proof.

For each $i \in \{2, \dots, m\}$, by Step 2, $V_i(\emptyset; f_i(R)) \geq -d_*$, by ia , $V_i(i; \mathbf{0}) = 4d_*$, and by Lemma 6 and Step 5, $t_i(R) \leq V_i(A_i(R); \mathbf{0}) = V_i(i; \mathbf{0})$. Thus, for each $i \in \{2, \dots, m\}$,

$$V_i(\emptyset; f_i(R)) \geq -d_* = V_i(i; \mathbf{0}) - 5d_* \geq t_i(R) - 5d_*. \quad (3)$$

By $A_1(R) = 1$, there is $A_1 \in \mathcal{M}$ such that $1 \in A_1$ and

$$V_1(A_1; f_1(R)) - t_1(R) = \max_{A'_1 \in \mathcal{M}_1} \{V_1(A'_1; f_1(R)) - t_1(R)\}.$$

By $A_1(R) = 1$, Lemma 6, and Lemma 9, $t_1(R) \in [0, V_1(1; \mathbf{0})]$. Thus, by the def. of \underline{t}_1 ,

$$V_1(A_1; f_1(R)) - t_1(R) \geq \underline{t}_1. \quad (4)$$

Let $N' \equiv \{i \in \{2, \dots, m\} : A_i(R) \subseteq A_1\}$. Let $z' \equiv ((A'_i, t'_i))_{i \in N} \in Z$ be such that

$$\begin{aligned} z'_1 &= (A_1, V_1(A_1; f_1(R))), \\ z'_i &= (\emptyset, V_i(\emptyset; f_i(R))) \text{ for each } i \in N', \text{ and} \\ z'_i &= f_i(R) \text{ for each } i \in N \setminus (N' \cup \{1\}). \end{aligned}$$

Figure 11 is an illustration of z' when $A_1 = \{1, 2, 3\}$.

***** FIGURE 11 (Illustration of z' when $A_1 = \{1, 2, 3\}$.) ENTERS HERE *****

Clearly, for each $i \in N$, $z'_i \leq f_i(R)$. By Step 5 and $1 \in A_1$, $|N'| = |A_1| - 1 \leq m - 1$. Thus,

$5|N'|d_* \leq 5(m-1)d_* < \underline{t}_1$, Therefore, by (3), (4), and $5|N'|d_* < \underline{t}_1$,

$$\begin{aligned}
\sum_{i \in N} t'_i &= V_1(A_1; f_1(R)) + \sum_{i \in N'} V_i(\emptyset; f_i(R)) + \sum_{i \in N \setminus (N' \cup \{1\})} t_i(R) \\
&\geq t_1(R) + \underline{t}_1 + \sum_{i \in N'} (t_i(R) - 5d_*) + \sum_{i \in N \setminus (N' \cup \{1\})} t_i(R) && \text{(by (3) and (4))} \\
&= \underline{t}_1 - 5|N'|d_* + \sum_{i \in N} t_i(R) \\
&> \sum_{i \in N} t_i(R). && \text{(by } 5|N'|d_* < \underline{t}_1)
\end{aligned}$$

This contradicts *efficiency*. ■

References

- [1] Adachi, Tsuyoshi (2014), “Equity and the Vickrey allocation rule on general preference domains.” *Social Choice and Welfare*, 42, 813–830.
- [2] Andersson, Tommy and Lars-Gunnar Svensson (2014), “Non-manipulable house allocation with rent control.” *Econometrica*, 82, 507–539.
- [3] Andersson, Tommy, Lars Ehlers, and Lars-Gunnar Svensson (2015), “Transferring ownership of public housing to existing tenants: A market design approach.” Working paper, Lund University.
- [4] Alkan, Ahmet and Gale, David (1990), “The core of the matching game.” *Games and Economic Behavior*, 2, 203–212.
- [5] Ashlagi, Itai and Shigehiro Serizawa (2012), “Characterizing Vickrey allocation rule by anonymity.” *Social Choice and Welfare*, 38, 531–542.
- [6] Ausubel, Lawrence M. (2004), “An efficient ascending-bid auction for multiple objects.” *American Economic Review*, 94, 1452–1475.
- [7] Ausubel, Lawrence M. (2006), “An efficient dynamic auction for heterogeneous commodities.” *American Economic Review*, 96, 602–629.
- [8] Ausubel, Lawrence M. and Paul R. Milgrom (2002), “Ascending auctions with package bidding.” *Frontiers of Theoretical Economics*, 1, 1–42.
- [9] Baisa, Brian (2013), “Auction design without quasilinear preferences.” Working paper, Amherst College.
- [10] Berga, Dolores and Shigehiro Serizawa (2000), “Maximal domain for strategy-proof rules with one public good.” *Journal of Economic Theory*, 90, 39–61.
- [11] Berge, Claude (1963), “Topological Spaces.” Macmillan, New York.

- [12] Bikhchandani, Sushil and Joseph M. Ostroy (2002), “The package assignment model.” *Journal of Economic Theory*, 107, 377–406.
- [13] Ching, Stephen and Shigehiro Serizawa (1998), “A maximal domain for the existence of strategy-proof rules.” *Journal of Economic Theory*, 78, 157–166.
- [14] Clarke, Edward H. (1971), “Multipart pricing of public goods.” *Public Choice*, 11, 17–33.
- [15] de Vries, Sven, James Schummer, and Rakesh V. Vohra (2007), “On ascending Vickrey auctions for heterogeneous objects.” *Journal of Economic Theory*, 132, 95–118.
- [16] Demange, Gabrielle and David Gale (1985), “The strategy structure of two-sided matching markets.” *Econometrica*, 53, 873–888.
- [17] Ehlers, Lars (2002), “Coalitional strategy-proof house allocation.” *Journal of Economic Theory*, 105, 298–317.
- [18] Gale, David (1984), “Equilibrium in a discrete exchange economy with money.” *International Journal of Game Theory*, 13, 61–64.
- [19] Gibbard, Allan (1973), “Manipulation of voting schemes: A general result.” *Econometrica*, 41, 587–601.
- [20] Groves, Theodore (1973), “Incentives in teams.” *Econometrica*, 41, 617–631.
- [21] Gul, Faruk and Ennio Stacchetti (1999), “Walrasian equilibrium with gross substitutes.” *Journal of Economic Theory*, 87, 95–124.
- [22] Gul, Faruk and Ennio Stacchetti (2000), “The English auction with differentiated commodities.” *Journal of Economic Theory*, 92, 66–95.
- [23] Holmström, Bengt (1979), “Groves’ scheme on restricted domains.” *Econometrica*, 47, 1137–1144.
- [24] Marshall, Alfred (1920), “Principles of economics.” Macmillan, London.
- [25] Massó, Jordi and Alejandro Neme (2001), “Maximal domain of preferences in the division problem.” *Games and Economic Behavior*, 37, 367–387.
- [26] Mishra, Debasis and David C. Parkes (2007), “Ascending price Vickrey auctions for general valuations.” *Journal of Economic Theory*, 132, 335–366.
- [27] Morimoto, Shuhei and Shigehiro Serizawa (2015), “Strategy-proofness and efficiency with non-quasi-linear preference: A characterization of minimum price Walrasian rule.” *Theoretical Economics*, 10, 445–487.
- [28] Myerson, Roger B. (1981), “Optimal auction design.” *Mathematics of Operations Research*, 6, 58–73.
- [29] Pápai, Szilvia (2003), “Groves sealed bid auctions of heterogeneous objects with fair prices.” *Social Choice and Welfare*, 20, 371–385.

- [30] Quinzii, Martine (1984), “Core and competitive equilibria with indivisibilities.” *International Journal of Game Theory*, 13, 41–60.
- [31] Sakai, Toyotaka (2008), “Second price auctions on general preference domains: Two characterizations.” *Economic Theory*, 37, 347–356.
- [32] Saitoh, Hiroki and Shigehiro Serizawa (2008), “Vickrey allocation rule with income effect.” *Economic Theory*, 35, 391–401.
- [33] Satterthwaite, Mark A. (1975), “Strategy-proofness and Arrow’s conditions: Existence and correspondence theorems for voting procedures and social welfare functions.” *Journal of Economic Theory*, 10, 187–217.
- [34] Sprumont, Yves (2013), “Constrained-optimal strategy-proof assignment: Beyond the Groves mechanisms.” *Journal of Economic Theory*, 148, 1102–1121.
- [35] Sun, Ning and Zaifu Yang (2006), “Equilibria and indivisibilities: Gross substitutes and complements.” *Econometrica*, 74, 1385–1402.
- [36] Sun, Ning and Zaifu Yang (2009), “Double-track adjustment process for discrete markets with substitutes and complements.” *Econometrica*, 77, 933–952.
- [37] Sun, Ning and Zaifu Yang (2014), “An efficient and incentive compatible dynamic auction for multiple complements.” *Journal of Political Economy*, 122, 422–466.
- [38] Tierney, Ryan (2015), “Managing multiple commons: Strategy-proofness and min-price Walras.” mimeo.
- [39] Vickrey, William. (1961), “Counterspeculation, auctions, and competitive sealed tenders.” *Journal of Finance*, 16, 8–37.




Figure 1: Unit-demand preference relation.





Figure 2: Multi-demand preference relation.






Figure 4: $R = (R_1, R_2, R_3)$.




Figure 5: (R'_1, R_{-1}) and z in Step C.




Figure 6: (R_1'', R_{-1}) and z' in Step C.




Figure 7: z in Step D.




Figure 9: Illustrations of R_i ($i \in \{2, \dots, m\}$) and R_j ($j \in \{m+1, \dots, m\}$).




Figure 10: Illustration of R_1'' .




Figure 11: Illustration of z' when $A_1 = \{1, 2, 3\}$.