

Identification Robust Testing of Risk Premia in Finite Samples

Frank Kleibergen* Lingwei Kong[†] Zhaoguo Zhan[‡]

June 17, 2019

Abstract

The reliability of tests on the risk premia in linear factor models is threatened by limited sample sizes and weak identification of risk premia frequently encountered in applied work. We propose novel tests on the risk premia that are robust to both limited sample sizes and the identification strength of the risk premia as reflected by the quality of the risk factors. These tests are appealing for empirically relevant settings, and lead to confidence sets of risk premia that can substantially differ from conventional ones. To show the latter, we revisit two high-profile empirical applications.

JEL Classification: G12

Keywords: asset pricing; risk premia; identification robust inference; finite samples

Acknowledgment: We thank [Paolo Zaffaroni, Imperial College Business School] and [Francisco Penaranda, QueensCollege CUNY] for their nice discussions that greatly improved the manuscript during Sofie 2019 in Shanghai.

*Amsterdam School of Economics, University of Amsterdam, Roetersstraat 11, 1018 WB Amsterdam, The Netherlands. Email: f.r.kleibergen@uva.nl.

[†]Amsterdam School of Economics, University of Amsterdam, Roetersstraat 11, 1018 WB Amsterdam, The Netherlands. Email: l.kong@uva.nl.

[‡]Department of Economics, Finance and Quantitative Analysis, Coles College of Business, Kennesaw State University, GA 30144, USA. Email: zzhan@kennesaw.edu.

1 Introduction

Over the past decades, it has become common practice to assess risk premia using the classic Fama-MacBeth (FM) two-pass approach, see *e.g.* Fama and MacBeth (1973) and Shanken (1992). It is based on the, so-called, linear beta representation of expected asset returns (Cochrane (2001)) and is one of the default approaches to evaluate asset pricing models in the financial economics literature, see *e.g.* Fama and French (1992). The large variety of risk factors that this literature has proposed (see *e.g.* Lettau and Ludvigson (2001), Kroencke (2017)) has, however, led to a growing concern that the reliability of the two-pass approach is at risk for many of them. This concern results from two common issues that threaten the two-pass approach: (i) the absence of large correlations between risk factors and asset returns needed to identify the risk premia; and (ii) the rather small number of time-series observations relative to the number of assets.

A strand of the literature shows a failure in identifying risk premia when the quality of risk factors specified for the two-pass approach is unsatisfactory. Kan and Zhang (1999), for instance, show that if risk factors are useless (so betas are zero), their risk premia are unidentified and the commonly used FM t -statistic on risk premia tends to be spuriously significant. Many observed risk factors are, however, not completely useless but just minorly correlated with asset returns. For these factors, Kleibergen (2009) similarly warns that their risk premia are only weakly identified, which also makes the FM t -statistic unreliable. These findings lead to the emergence of the so-called identification robust tests on risk premia, which, unlike the FM t -test, remain trustworthy regardless of the quality of risk factors. In the ideal scenario where risk factors of good quality are used for the FM two-pass approach, identification robust tests and the FM t -test both provide valid and comparable inference on risk premia. In contrast, in empirical studies where the quality of risk factors is questionable, identification robust tests are more trustworthy than the FM t -test. See, *e.g.*, Kleibergen (2009), Beaulieu, Dufour and Khalaf (2013), Khalaf and Schaller (2016), Kleibergen and Zhan (2018).

Another strand of literature is concerned with the challenges that result from the limited

sample sizes in empirical studies. Macroeconomic risk factors, in particular, are commonly measured at quarterly or annual frequencies, so their number of time-series observations T is usually not much larger than the number of test assets N . Take Kroencke (2017), for example, the yearly consumption growth series from 1960–2014 has fifty-five time-series observations, $T = 55$, while thirty portfolios, sorted by size, value and investment alongside the market portfolio, are used as test assets, so $N = 31$. This limited T vs. large N setting jeopardizes asymptotic tests on the risk premia, whose validity typically relies on a much larger value of T compared to N (Newey and Windmeijer (2009)). To conduct valid inference for limited sample sizes, researchers have therefore developed various finite-sample robust methods. Gibbons, Ross and Shanken (GRS) (1989), for example, propose the GRS test for portfolio efficiency, which circumvents the limit sample size problem by making distributional assumptions. Similar assumptions are made in Kan and Zhou (2004) for deriving the exact distribution of the Hansen-Jagannathan statistic. More recently, Kleibergen and Zhan (2019) extend the GRS test to develop a finite-sample robust test on the risk premia.

Both strands of literature above are empirically relevant, since sample sizes in empirical studies are often not sufficiently large to validate asymptotic tests, such as the FM t -test, while it has been well documented that a large number of risk factors are of questionable quality (Kleibergen and Zhan (2015)). Despite the fact that the above two strands of literature are well established, tests on risk premia that account for both the limited sample size and the questionable quality of risk factors are lacking.

In light of the above, we propose novel tests on the risk premia, which are designed to be robust to both the sample size and the quality of risk factors. We do so by providing the exact finite-sample distributions of several identification robust test statistics, including those in, e.g., Kleibergen (2009) and Beaulieu, Dufour and Khalaf (2013). Put differently, while identification robust tests are asymptotically valid regardless of the quality of risk factors, we extend these tests to finite-sample settings by constructing finite-sample distributions of their test statistics. In line with Gibbons, Ross and Shanken (1989), Kleibergen and Zhan (2019) and many others, these finite-sample distributions result from the (normality)

assumptions of the error term in a linear factor model for asset returns.

We show that our proposed tests on risk premia work favorably in a simulation experiment calibrated to real data. As expected, these tests are found particularly appealing in simulation settings where the time-series dimension T is close to the cross-section dimension N . In such settings, the proposed tests lead to correct sizes for testing hypotheses on risk premia, while their asymptotic counterparts, i.e., the asymptotically valid tests, can suffer from size distortion. On the other hand, when T is much larger than N , the proposed tests become identical to the asymptotically valid identification robust tests, all of which are trustworthy regardless of the quality of risk factors. These numerical findings for the proposed tests are all in line with our theoretical results.

To illustrate the practical relevance of our proposed tests, we revisit the empirical studies on the consumption Capital Asset Pricing Model (CAPM) and the conditional consumption CAPM in Kroencke (2017) and Lettau and Ludvigson (2001), respectively.¹ Kroencke (2017) focuses on the classic consumption CAPM, where consumption growth is the single risk factor. The involved data set has $T = 55$ while $N = 31$ as aforementioned. Instead of the single factor model, Lettau and Ludvigson (2001) propose a multi-factor conditional consumption CAPM. We use their quarterly data from 1963Q3-1998Q3 with $T = 141$ and $N = 25$ Fama-French portfolios sorted by size and book-to-market as test assets. In both applications, we find that inverting the proposed tests leads to 95% confidence sets of risk premia that are substantially different from those based on existing asymptotic tests. This should not be surprising, since both in Lettau and Ludvigson (2001) and Kroencke (2017), the limited T relative to N is unlikely to validate asymptotic tests. Moreover, the 95% confidence sets resulting from the proposed tests are found to be unbounded, reflecting that little information on risk premia is contained in the studied data sets. These empirical results thus cast doubt on the pricing performance of the proposed risk factors in Lettau and Ludvigson (2001) and Kroencke (2017).

Overall, our findings indicate that robustness to the quality of risk factors and robustness

¹We thank these authors for providing their data.

to the sample size should both be accounted for in asset pricing studies. Traditional asset pricing tests that are not robust to the strength of identification reflected by the quality of risk factors can lead to erroneous conclusions. Similarly, asymptotically valid tests that do not account for limited sample sizes can also induce spurious empirical findings. We simultaneously address these two empirically relevant issues.

The rest of the paper is organized as follows. Section 2 discusses the two issues that threaten reliable inference on risk premia: lack of identification due to poor quality risk factors and the limited sample size. It also introduces the two data sets from Kroencke (2017) and Lettau and Ludvigson (2001) that we use to emphasize the empirical relevance of our results. Section 3 presents the proposed test statistics and constructs their finite-sample distributions. Section 4 contains a simulation study as well as applications using the data from Lettau and Ludvigson (2001) and Kroencke (2017). Section 5 concludes. Technical details are relegated to the Appendix.

2 Preliminaries

2.1 Linear beta representation and the two-pass approach

Premia on risk factors are identified by the linear beta representation of expected asset returns (Cochrane (2001)):

$$\mathbb{E}(R_t) = \iota_N \lambda_0 + \beta \lambda_F = (\iota_N, \beta) \begin{pmatrix} \lambda_0 \\ \lambda_F \end{pmatrix}, \quad (1)$$

with R_t an $N \times 1$ vector of asset returns, ι_N the $N \times 1$ vector of ones, β an $N \times K$ matrix of factor loadings, λ_0 the zero-beta return, and λ_F the $K \times 1$ vector of risk premia.

The β matrix of factor loadings in (1) is related to a linear factor model for asset returns:

$$R_t = c + \beta F_t + u_t, \quad t = 1, \dots, T, \quad (2)$$

with F_t a $K \times 1$ vector of risk factors, c an $N \times 1$ vector of constant terms, and u_t an $N \times 1$ vector of errors.

A large portion of the asset pricing literature is centered around inference on the risk premia λ_F based on (1)-(2). The conventional two-pass procedure estimates β in the first pass using the time-series regression in (2), so

$$\hat{\beta} = \left(\sum_{t=1}^T \bar{R}_t \bar{F}_t' \right) \left(\sum_{t=1}^T \bar{F}_t \bar{F}_t' \right)^{-1}, \quad (3)$$

with $\bar{R}_t = R_t - \bar{R}$, $\bar{R} = \frac{1}{T} \sum_{t=1}^T R_t$, $\bar{F}_t = F_t - \bar{F}$, $\bar{F} = \frac{1}{T} \sum_{t=1}^T F_t$; in the second pass, the risk premia are estimated using a cross-sectional regression of the average asset returns \bar{R} on ι_N and $\hat{\beta}$ estimated in the first pass:

$$\begin{pmatrix} \hat{\lambda}_0 \\ \hat{\lambda}_F \end{pmatrix} = \left[\left(\iota_N, \hat{\beta} \right)' \left(\iota_N, \hat{\beta} \right) \right]^{-1} \left(\iota_N, \hat{\beta} \right)' \bar{R}. \quad (4)$$

2.2 Lack of identification and identification robust tests

The (ι_N, β) matrix plays a crucial role for inference on risk premia (Kleibergen and Zhan (2019)). If (ι_N, β) has a full rank value, then λ_F is well defined by (1), and its estimator $\hat{\lambda}_F$ is consistent and asymptotically normally distributed under standard regularity conditions (Shanken (1992)). It validates the common usage of the t -test on the risk premia. The full rank condition of (ι_N, β) is, however, plausibly violated in many empirical settings.

The pioneering study of Kan and Zhang (1999), for example, focuses on the useless factor setting with $\beta = 0$. If $\beta = 0$, then λ_F is unidentified in (1), and Kan and Zhang (1999) show that the FM t -test on risk premia becomes unreliable, i.e., its test statistic tends to be significant, which thus spuriously supports useless risk factors. Kleibergen (2009) further considers a so-called weak factor setting with $\beta \approx 0$, since many risk factors proposed in the literature exhibit just minor correlation with asset returns. Similar to the useless factor setting, Kleibergen (2009) finds that risk premia on weak factors are not well identified,

which further induces a malfunction of the FM t -test. In addition, Kleibergen and Zhan (2019) show that β 's are often close to be constant over the different assets so there is little cross-sectional variation in them. Kleibergen and Zhan (2019) warn that identification of the risk premia is then similarly jeopardized, so the FM t -test also breaks down. All these studies boil down to the full rank condition of (ι_N, β) , which is jeopardized if $\beta = 0$, $\beta \approx 0$ or $\beta \propto \iota_N$.

To provide empirical evidence that the full rank condition is plausibly violated, we use data from Kroencke (2017) and Lettau and Ludvigson (2001). The resulting estimates of β are presented in Table 1 and Table 2, respectively.

Re-produced from Kleibergen and Zhan (2019), Table 1 presents the estimated β 's using the thirty-one test assets from Kroencke (2017) and five different consumption growth measures (“Reported”, “P-J”, “Q4-Q4”, “Garbage”, “Unfiltered”, see the notes to Table 1) from the asset pricing literature. Kroencke (2017) proposes “Unfiltered” consumption growth as a risk factor, which appears to outperform the other four consumption growth measures for consumption-based asset pricing.

Table 1, however, suggests that the β 's for all five consumption growth measures are problematic. For example, the estimated β -vector for “Reported” consumption growth is not statistically different from zero. On the other hand, although the β 's for the “Garbage” and “Unfiltered” consumption growth measures all appear sizeable, they exhibit little cross-sectional variation so they are almost constant and proportional to a vector of ones. Kleibergen and Zhan (2019) further conduct a rank test on (ι_N, β) and find evidence that its full rank condition is likely violated under each of the five consumption growth measures. The reported estimates of the β 's in Table 1 thus cast doubt on the empirical findings in Kroencke (2017).

Similar to Table 1, Table 2 reports the estimated β 's using the twenty-five Fama-French portfolios sorted by size and book-to-market for three risk factors (the quarterly consumption growth Δc , the lagged consumption-wealth ratio cay and their interaction $\Delta c \times cay$) taken from Lettau and Ludvigson (2001). It is rather obvious in Table 2 that the last column of the β -matrix, which corresponds to the $\Delta c \times cay$ risk factor, is not statistically significant

Table 1: β with thirty-one portfolio returns

	Reported		P-J		Q4-Q4		Garbage		Unfiltered	
	$\hat{\beta}$	t -stat	$\hat{\beta}$	t -stat	$\hat{\beta}$	t -stat	$\hat{\beta}$	t -stat	$\hat{\beta}$	t -stat
(1)	-1.30	-0.41	1.19	0.93	4.18	1.51	4.54	3.33	5.40	3.81
(2)	-0.23	-0.09	1.14	1.06	4.42	1.93	4.06	3.52	4.70	3.98
(3)	0.14	0.06	0.99	1.06	4.15	2.09	3.74	3.86	3.80	3.64
(4)	-0.52	-0.25	0.73	0.86	3.46	1.92	3.66	4.30	3.33	3.50
(5)	-0.23	-0.11	0.73	0.85	3.74	2.06	3.62	4.30	3.33	3.44
(6)	-0.84	-0.43	0.58	0.72	3.01	1.75	3.42	4.39	3.10	3.43
(7)	-0.85	-0.44	0.49	0.63	2.73	1.63	3.51	4.90	2.95	3.32
(8)	-0.59	-0.33	0.39	0.53	2.69	1.69	3.06	4.55	2.85	3.41
(9)	-0.12	-0.07	0.30	0.43	2.75	1.87	2.61	3.98	2.48	3.16
(10)	0.34	0.20	0.64	0.92	2.97	2.01	3.14	4.53	2.60	3.29
(11)	-0.95	-0.46	0.21	0.25	2.25	1.22	3.58	4.29	2.85	2.91
(12)	-0.44	-0.25	0.60	0.81	2.77	1.77	3.15	4.22	2.59	3.09
(13)	-0.63	-0.38	0.52	0.78	2.49	1.73	2.95	4.38	2.22	2.84
(14)	-0.56	-0.31	0.07	0.10	2.26	1.39	2.51	3.10	2.04	2.28
(15)	0.88	0.53	0.46	0.67	3.14	2.19	2.75	3.89	2.25	2.85
(16)	0.73	0.45	0.58	0.88	3.04	2.19	2.86	4.49	2.62	3.56
(17)	0.96	0.56	0.71	1.03	3.50	2.40	2.93	4.53	2.81	3.61
(18)	1.21	0.65	1.05	1.40	4.27	2.72	3.17	4.35	3.33	4.02
(19)	1.23	0.67	1.09	1.47	3.97	2.56	2.86	3.65	2.96	3.53
(20)	1.77	0.84	1.32	1.56	5.22	2.98	2.93	3.26	4.21	4.67
(21)	-0.50	-0.22	0.58	0.61	3.57	1.78	3.85	4.30	3.88	3.73
(22)	-0.09	-0.05	0.66	0.88	3.14	1.98	3.00	4.10	3.15	3.82
(23)	0.37	0.23	0.54	0.81	2.84	2.02	2.94	4.49	2.27	2.96
(24)	0.44	0.31	0.54	0.92	2.96	2.40	2.84	5.30	2.25	3.38
(25)	-0.50	-0.32	0.31	0.49	2.11	1.57	2.69	5.05	2.22	3.10
(26)	1.04	0.62	0.79	1.16	3.70	2.60	3.09	4.44	2.54	3.25
(27)	1.10	0.66	0.98	1.46	3.71	2.63	2.90	4.13	2.74	3.60
(28)	0.03	0.01	0.71	0.98	3.18	2.07	3.15	4.32	3.07	3.83
(29)	-0.89	-0.41	0.17	0.19	2.44	1.26	3.54	3.82	2.70	2.59
(30)	-1.40	-0.58	0.03	0.03	2.10	0.97	3.92	3.95	2.92	2.52
(31)	0.23	0.13	0.72	1.00	3.32	2.21	3.21	4.74	2.95	3.72

Notes: The thirty-one test assets are taken from Kroencke (2017), i.e., the thirty portfolios sorted by size, value and investment, plus the market portfolio in 1960-2014 (yearly data). “Reported”, “P-J”, “Q4-Q4”, “Garbage” and “Unfiltered” correspond to five consumption measures: (i) the consumption expenditure on nondurable goods and services reported by National Income and Product Accounts (NIPA) (“Reported”); (ii) the three-year consumption measure from Parker and Julliard (2005) (“P-J”); (iii) the fourth-quarter to fourth-quarter consumption measure from Jagannathan and Wang (2007) (“Q4-Q4”); (iv) the garbage measure in Savov (2011) (“Garbage”); (v) the unfiltered NIPA consumption measure in Kroencke (2017) (“Unfiltered”). Each of the five consumption growth rates is used as the single risk factor for asset returns to estimate β .

Table 2: β with twenty-five portfolio returns

	Δc		cay		$\Delta c \times \mathbf{cay}$	
	$\hat{\beta}_1$	<i>t</i> -stat	$\hat{\beta}_2$	<i>t</i> -stat	$\hat{\beta}_3$	<i>t</i> -stat
(1)	6.35	2.32	4.25	2.96	-4.56	-0.22
(2)	6.30	2.61	3.66	2.90	2.25	0.12
(3)	5.11	2.28	3.22	2.75	3.22	0.19
(4)	5.50	2.58	2.96	2.65	5.12	0.32
(5)	5.79	2.53	2.65	2.21	11.36	0.65
(6)	4.38	1.76	4.55	3.49	-15.82	-0.84
(7)	3.63	1.65	3.25	2.83	0.02	0.00
(8)	3.92	1.97	3.07	2.95	-1.08	-0.07
(9)	3.52	1.91	2.66	2.76	6.93	0.50
(10)	4.83	2.41	2.16	2.06	8.48	0.56
(11)	2.70	1.21	4.53	3.88	-22.20	-1.31
(12)	2.76	1.47	3.64	3.69	-3.35	-0.23
(13)	2.92	1.69	2.63	2.91	4.40	0.34
(14)	2.58	1.57	2.73	3.18	0.02	0.00
(15)	3.71	1.98	2.15	2.19	4.78	0.34
(16)	1.97	1.02	4.22	4.20	-20.28	-1.39
(17)	2.62	1.49	3.28	3.58	-10.24	-0.77
(18)	1.94	1.21	2.57	3.06	-4.98	-0.41
(19)	2.50	1.56	2.32	2.76	0.60	0.05
(20)	3.78	2.05	2.26	2.34	3.28	0.23
(21)	1.61	1.01	2.77	3.33	-21.22	-1.76
(22)	1.16	0.80	2.53	3.33	-4.20	-0.38
(23)	2.32	1.88	2.46	3.80	-8.27	-0.88
(24)	1.24	0.94	2.24	3.25	-10.49	-1.05
(25)	3.07	2.13	1.59	2.11	-1.25	-0.11

Notes: The twenty-five Fama-French portfolios sorted by size and book-to-market in 1963Q3-1998Q3 (quarterly data) are used as test assets. The three risk factors taken from Lettau and Ludvigson (2001) are the quarterly consumption growth Δc , the (lagged) consumption-wealth ratio \mathbf{cay} and their interaction $\Delta c \times \mathbf{cay}$. The 25×3 dimensional β matrix is thus $\beta = (\beta_1, \beta_2, \beta_3)$, where β_i , $i = 1, 2, 3$, corresponds to the i -th factor.

at the 5% level. We therefore can not rule out that a column of the β -matrix for this multi-factor model equals zero, so the (ι_N, β) matrix does not have a full rank value. This is also consistent with the rank test results reported in Kleibergen and Zhan (2019). Consequently, Table 2 raises doubts on the credibility of the empirical findings based on the FM t -test in Lettau and Ludvigson (2001).

The β 's of questionable quality, such as those reported in Tables 1-2, have motivated Kan and Zhang (1999), Kleibergen (2009) and thereafter Kleibergen and Zhan (2015, 2018, 2019), to analyze their impact on the risk premia. It is now well acknowledged that inference on the risk premia is sensitive to the strength of identification, as reflected by the full rank condition of (ι_N, β) or put differently, the quality of the risk factors that yield the factor loadings in β . This has led to the so-called identification robust tests on the risk premia, whose validity does not depend on the rank of (ι_N, β) , i.e., these tests remain trustworthy regardless of the quality of the risk factors. An example of such a test is the factor Anderson-Rubin (FAR) test proposed by Kleibergen (2009), which is an extension of the robust Anderson and Rubin (1949) test in the instrumental variables regression literature, see also the Hotelling type test in Beaulieu, Dufour and Khalaf (2013). Other examples of identification robust tests include the Lagrange multiplier (LM) and likelihood ratio (LR) tests.

One potential drawback of these identification robust tests is, however, that many of them are established using asymptotic arguments, so the validity of these tests often relies on a large sample size, see, e.g., the FAR test in Kleibergen (2009). It is thus questionable whether these tests perform well in finite-sample applications, which we discuss next.

2.3 Limited T vs. large N and finite-sample robust tests

Other than lack of identification due to the quality of risk factors, limited sample sizes also impose challenges for inference on risk premia. The latter should come out naturally, since validity of asymptotic tests, such as the FM t -test and the FAR test, requires a large number of time-series observations T relative to the number of assets N . If T is small while N is large, it becomes doubtful whether we can still approximate the distributions of the FM

t -statistic and the FAR statistic by normal and chi-squared distributions, respectively. The large number of assets N or in other words, many moment conditions, resembles the many instruments issue in instrumental variables regression (Bekker and Kleibergen (2003), Bekker (1994), Newey and Windmeijer (2009)).

The limited T vs. large N scenario is commonly encountered in empirical asset pricing studies. For instance, Savov (2011) uses the annual garbage growth series from 1960–2006, while the twenty-five Fama-French portfolios augmented by the ten industry portfolios, as suggested by Lewellen, Nagel and Shanken (2010), are used as test assets. The resulting limited $T = 47$ and relatively large $N = 35$ thus cast doubt on the FM t -test used in Savov (2011). Similarly, the Kroencke (2017) data used for Table 1 uses the yearly unfiltered consumption growth series for the time period 1960–2014, so $T = 55$, and thirty portfolios sorted by size, value and investment alongside the market portfolio are used as test assets, so $N = 31$. The simulation experiment in Kleibergen and Zhan (2019), however, shows that neither the FM t -test nor the FAR test is size-correct under the limited T vs. large N setting calibrated to Kroencke (2017). Given that many risk factors, like the garbage growth in Savov (2011) and consumption growth in Kroencke (2017), are only available at low frequencies and/or within short time periods, the limited T vs. large N problem similarly affects many other empirical studies.

In view of the above, we aim to develop tests on risk premia that are immune to the limited T vs. large N problem, and we label these tests as finite-sample robust tests. Meanwhile, we also want these tests to remain trustworthy when the quality of risk factors is potentially unsatisfactory. We therefore have a two-fold goal, i.e., providing tests that are reliable when using risk factors of questionable quality as well as for limited sample sizes.

Kleibergen and Zhan (2019) develop two tests to meet this two-fold goal. First, they construct a finite-sample version of the rank test from Kleibergen and Paap (2006) so it can be used to test for a reduced rank value of (ι_N, β) , implicating that the risk premia are not identified, when N is large compared to T . Second, they extend the GRS test statistic from Gibbons, Ross and Shanken (1989) so that it can be used for testing hypotheses on the risk

premia. They refer to it as the GRS-FAR statistic, to reflect that it also results from the FAR statistic in Kleibergen (2009) with an adjustment for the finite-sample setting. As in Gibbons, Ross and Shanken (1989), Kleibergen and Zhan (2019) also assume that the error term in the linear factor model (2) is normally distributed, in order to derive the finite-sample F -distribution of the GRS-FAR statistic. Since this F -distribution does not depend on the quality of the risk factors, the finite-sample robust GRS-FAR test is also robust to the strength of identification.

In the next section, we propose several other tests, which, like the GRS-FAR test, are robust to both the sample size and the strength of identification reflected by the quality of the risk factors. Later on, we also show that these tests have comparable or even better power than the GRS-FAR test. To facilitate comparison and understanding, we also discuss the GRS-FAR test.

3 Analytical results

3.1 Null hypothesis and assumptions

Our interest lies in testing the hypothesis on the risk premia: $H_0 : \lambda_F = \lambda_{F,0}$, with $\lambda_{F,0}$ the hypothesized value of λ_F . We therefore delete λ_0 in (1) by removing the N -th asset and taking all other asset returns in deviation from the return on the N -th asset. The moment condition (1) and the linear factor model (2) now become:

$$\mathbb{E}(\mathcal{R}_t) = \mathcal{B}\lambda_F \tag{5}$$

$$\mathcal{R}_t = \mathcal{C} + \mathcal{B}F_t + \mathcal{U}_t \tag{6}$$

where $\mathcal{R}_t = J_N R_t$, $\mathcal{C} = J_N c$, $\mathcal{B} = J_N \beta$, $\mathcal{U}_t = J_N u_t$ with $J_N = (I_{N-1}, -\iota_{N-1})$, I_{N-1} is the $(N-1) \times (N-1)$ identity matrix, and ι_{N-1} is the $(N-1)$ -dimensional vector of ones.

Remark 1. Our proposed test statistics are invariant to which asset return is to be subtracted. Put differently, the choice of the N -th asset does not affect the resulting values

of our statistics. See the Internet Appendix of Kleibergen and Zhan (2019).

Remark 2. In tests with excess returns, $\lambda_0 = 0$ is sometimes imposed. See, e.g., Savov (2011). With this zero-restriction, we do not have to remove the N -th asset, and (5)-(6) are almost identical to (1)-(2) except for notation. Our tests can thus be extended straightforwardly to incorporate the $\lambda_0 = 0$ restriction.

Remark 3. The full rank condition of (ι_N, β) , which we discussed previously for identifying risk premia, is equivalent to the full rank condition of \mathcal{B} . Since we allow for risk factors of questionable quality, we do not need to assume that \mathcal{B} has a full rank value.

We make the following two assumptions for the linear factor model, both of which have been used in the previous literature. Further explanations are provided under each assumption.

Assumption 1. \mathcal{U}_t is *i.i.d.* normally distributed with covariance Σ .

The normality assumption gives rise to the finite-sample distributions of the test statistics proposed later on. This assumption is commonly adopted in the existing literature to facilitate finite-sample analysis, see, e.g., Gibbons, Ross and Shanken (1989), Kan and Zhou (2004). In addition, there also exists empirical evidence that distributions of some asset returns could be approximated by the normal distribution (e.g., Kleibergen and Zhan (2019)). We can as well replace normality with other distributional assumptions, and if we do so, the finite-sample distributions of our proposed statistics can be derived in a similar fashion or using simulation.

Assumption 2. Let F be the $T \times K$ matrix whose t -th row is \bar{F}_t' , and we assume that (ι_T, F) is a deterministic matrix of full rank.

The full rank condition of (ι_T, F) , where ι_T is the T -dimensional vector of ones, rules out multi-collinearity of risk factors, so β and thus \mathcal{B} are estimable using time-series regression. Here we assume that risk factors are fixed, instead of imposing restrictions on their limiting behaviors, in order to facilitate the finite-sample analysis of test statistics. The assumption of fixed factors can be similarly found in, e.g., Gibbons, Ross and Shanken (1989). We could

allow for randomness of risk factors by regarding the proposed test statistics as conditional random variables given the risk factors.

3.2 Test statistics and their finite-sample distributions

Under the null hypothesis $H_0 : \lambda_F = \lambda_{F,0}$ and the moment condition (5), we have $\mathbb{E}(\mathcal{R}_t) = \mathcal{B}\lambda_{F,0}$, so the average returns $\bar{\mathcal{R}} = \frac{1}{T} \sum_{t=1}^T \mathcal{R}_t$ should be centered around $\mathcal{B}\lambda_{F,0}$. In order to test $H_0 : \lambda_F = \lambda_{F,0}$, we can examine whether $\bar{\mathcal{R}} - \mathcal{B}\lambda_{F,0}$ is centered around zero. This motivates the test statistics we present next.

Put differently, our proposed test statistics do not rely on an estimator of λ_F . Instead, we just examine whether the factor pricing condition (5) holds at the hypothesized risk premia $\lambda_{F,0}$. A rejection of $H_0 : \lambda_F = \lambda_{F,0}$ could result if (5) holds at a value other than $\lambda_{F,0}$, or if (5) does not hold at any value of the risk premia, i.e., there exists model mis-specification.² Since our test statistics do not involve risk premia estimation, they remain trustworthy regardless of the strength of risk premia identification, as reflected by the quality of risk factors or \mathcal{B} .

For later use, we distinguish the following two estimators for \mathcal{B} : a restricted least squares estimator under the null,

$$\tilde{\mathcal{B}} = \left(\sum_{t=1}^T \mathcal{R}_t (\bar{F}_t + \lambda_{F,0})' \right) \left(\sum_{t=1}^T (\bar{F}_t + \lambda_{F,0}) (\bar{F}_t + \lambda_{F,0})' \right)^{-1},$$

and an unrestricted least squares estimator,

$$\hat{\mathcal{B}} = \left(\sum_{t=1}^T \bar{\mathcal{R}}_t \bar{F}_t' \right) \left(\sum_{t=1}^T \bar{F}_t \bar{F}_t' \right)^{-1},$$

where $\bar{\mathcal{R}}_t = \mathcal{R}_t - \bar{\mathcal{R}}$, \bar{F}_t is the demeaned factor similarly defined under (3).

Since $\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0}$ and $\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}$ are natural choices to help examine whether $\mathbb{E}(\mathcal{R}_t) = \mathcal{B}\lambda_{F,0}$ holds, we use them to construct test statistics. Under $H_0 : \lambda_F = \lambda_{F,0}$ imposed on factor

²If $H_0 : \lambda_F = \lambda_{F,0}$ is rejected for every hypothesized $\lambda_{F,0}$, then it signals that the adopted risk factors are mis-specified for the factor pricing condition (5). In this case, mis-specification robust inference methods can be adopted (see, e.g., Kan, Robotti and Shanken (2013)).

pricing and Assumptions 1-2, $\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0}$ and $\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}$ are both normally distributed $(N-1)$ -dimensional vectors, and the distribution of one of these two vectors is just a scaled version of the distribution of the other:³

$$\sqrt{T} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right) \sim N \left(0, \left(1 - \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} \lambda_{F,0} \right) \otimes \Sigma \right), \quad (7)$$

$$\sqrt{T} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0} \right) \sim N \left(0, \left(1 + \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} \lambda_{F,0} \right) \otimes \Sigma \right), \quad (8)$$

where $\hat{\mathcal{Q}}_{FF}(\lambda_{F,0}) = \frac{1}{T} \sum_{t=1}^T (\bar{F}_t + \lambda_{F,0})(\bar{F}_t + \lambda_{F,0})'$, $\hat{\mathcal{Q}} = \frac{1}{T} \sum_{t=1}^T \bar{F}_t \bar{F}_t'$, and Σ is the covariance of the error \mathcal{U}_t in (6).

3.2.1 FAR and Hotelling (H) statistics

The FAR statistic in Kleibergen (2009) is based on the normal approximation of the $\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0}$ vector. Similarly, the Hotelling (H) type test statistic (see Beaulieu, Dufour and Khalaf (2013)) is based on the normal approximation of $\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}$. Both statistics can then be viewed as the standardized lengths of these vectors, which explains their χ^2 limiting distributions:

$$FAR(\lambda_{F,0}) = \frac{T}{1 - \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right)' \hat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right) \quad (9)$$

$$H(\lambda_{F,0}) = \frac{T}{1 + \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} \lambda_{F,0}} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0} \right)' \hat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0} \right) \quad (10)$$

with $\hat{\Sigma}$ a consistent estimator for the covariance matrix Σ . Unless specified otherwise, we use

$$\hat{\Sigma} = \frac{1}{T - K - 1} \sum_{t=1}^T \left(\bar{\mathcal{R}}_t - \hat{\mathcal{B}}\bar{F}_t \right) \left(\bar{\mathcal{R}}_t - \hat{\mathcal{B}}\bar{F}_t \right)'$$

as the covariance matrix estimator of Σ for the proposed test statistics.

Theorem 1. *If the same covariance estimator $\hat{\Sigma}$ is used for the FAR and Hotelling statistics, then $FAR(\lambda_{F,0}) = H(\lambda_{F,0})$. Under $H_0 : \lambda_F = \lambda_{F,0}$ and Assumptions 1-2, the FAR and H*

³See Lemmas 1 and 2 in the Appendix. For ease of exposition, all proofs are provided in the Appendix, so we can focus on our prime results in the main text.

statistics converge to a χ_{N-1}^2 distributed random variable, as $T \rightarrow \infty$ while N is fixed.

Proof: See the Appendix.⁴

Kleibergen and Zhan (2019) provide another interpretation of the FAR statistic by relating it to the well-known GRS statistic. They show that $\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0}$ can be viewed as an estimator of the constant terms (alphas) in a set of appropriately specified regressions of asset returns on risk factors. While alphas are zero under the null, the FAR statistic, like the GRS statistic, just tests whether alphas are zero. Therefore, we could also label the FAR statistic as the GRS-FAR statistic, as in Kleibergen and Zhan (2019).

The asymptotic result in Theorem 1, however, is at risk in the finite-sample setting, where the time-series dimension T may not be much larger than the cross-section dimension N . We therefore adjust the FAR statistic (or equivalently, the H statistic) and derive its finite-sample distribution, which is provided in Theorem 2.

Theorem 2. *Under $H_0 : \lambda_F = \lambda_{F,0}$ and Assumptions 1-2,*

$$\frac{T - K - N + 1}{(T - K - 1)(N - 1)} \times FAR(\lambda_{F,0}) \sim F_{N-1, T-K-N+1} \quad (11)$$

where $F_{N-1, T-K-N+1}$ is the F -distribution with $N - 1$, $T - K - N + 1$ degrees of freedom.

Proof. See the Appendix.

The finite-sample F -distribution in Theorem 2 indicates the importance of the time-series dimension T relative to the cross-section dimension N . If T is much larger than N , then it is straightforward to verify that the finite-sample result in Theorem 2 is equivalent to the asymptotic result in Theorem 1. On the other hand, if T is close to N , which is not uncommon in practice, then inference on risk premia based on Theorem 2 is expected to be substantially different from that based on Theorem 1. The related simulation evidence can be found in Kleibergen and Zhan (2019).

⁴Theorem 1, as well as other asymptotic results presented later on, also holds under weaker conditions, see Kleibergen (2009).

3.2.2 GLS-LM and JGLS statistics

As stated in Theorem 1, asymptotically the FAR statistic has a χ^2 distribution with $N - 1$ degrees of freedom. Therefore, when we use a large number of test assets, so the degrees of freedom increases, the FAR test becomes less powerful. This issue has been similarly discussed for the Anderson and Rubin (1949) statistic in the instrumental variables regression literature (Kleibergen (2002)).

To reduce the degrees of freedom and thus improve the power of tests on the risk premia, Kleibergen (2009) decomposes the FAR statistic into two parts, the GLS-LM statistic and the JGLS statistic, both can be used for testing $H_0 : \lambda_F = \lambda_{F,0}$:

$$FAR(\lambda_{F,0}) = GLS-LM(\lambda_{F,0}) + JGLS(\lambda_{F,0}).$$

The GLS-LM statistic results from projecting $\widehat{\Sigma}^{-\frac{1}{2}} (\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0})$ on the column space spanned by $\widehat{\Sigma}^{-\frac{1}{2}}\tilde{\mathcal{B}}$, so

$$\begin{aligned} & GLS-LM(\lambda_{F,0}) \\ &= \frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} (\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0})' \widehat{\Sigma}^{-\frac{1}{2}'} P_{\widehat{\Sigma}^{-\frac{1}{2}}\tilde{\mathcal{B}}} \widehat{\Sigma}^{-\frac{1}{2}} (\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0}) \\ &= \frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} (\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0})' \widehat{\Sigma}^{-1} \tilde{\mathcal{B}} (\tilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \tilde{\mathcal{B}})^{-1} \tilde{\mathcal{B}}' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0}) \end{aligned} \quad (12)$$

where $P_{\widehat{\Sigma}^{-\frac{1}{2}}\tilde{\mathcal{B}}} = \widehat{\Sigma}^{-\frac{1}{2}} \tilde{\mathcal{B}} (\tilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \tilde{\mathcal{B}})^{-1} \tilde{\mathcal{B}}' \widehat{\Sigma}^{-\frac{1}{2}'}$, $\widehat{\Sigma}^{-1} = \widehat{\Sigma}^{-\frac{1}{2}'} \widehat{\Sigma}^{-\frac{1}{2}}$. Moreover, the GLS-LM statistic can also be viewed as a quadratic form of the derivative of the FAR statistic (Kleibergen (2009)).

Similarly, the JGLS statistic results from projecting $\widehat{\Sigma}^{-\frac{1}{2}} (\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0})$ on the space orthogonal to the one spanned by $\widehat{\Sigma}^{-\frac{1}{2}}\tilde{\mathcal{B}}$, so it equals $FAR(\lambda_{F,0}) - GLS-LM(\lambda_{F,0})$:

$$\begin{aligned}
& JGLS(\lambda_{F,0}) \\
&= \frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \right)' \widehat{\Sigma}^{-\frac{1}{2}} (I_{N-1} - P_{\widehat{\Sigma}^{-\frac{1}{2}} \tilde{\mathcal{B}}}) \widehat{\Sigma}^{-\frac{1}{2}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \right) \quad (13)
\end{aligned}$$

By decomposing the FAR statistic as described above, the distributions of the resulting GLS-LM and JGLS statistics have fewer degrees of freedom, as shown in Theorem 3.

Theorem 3. *Under $H_0 : \lambda_F = \lambda_{F,0}$ and Assumptions 1-2, the GLS-LM and JGLS statistics converge to independent χ_K^2 and χ_{N-K-1}^2 distributed random variables, respectively, as $T \rightarrow \infty$ while N is fixed.*

Proof: See the Appendix.

To make GLS-LM and JGLS tests applicable for the limited T vs. large N setting, we similarly derive the finite-sample distributions of their (re-scaled) statistics. These are provided in Theorem 4.

Theorem 4. *Under $H_0 : \lambda_F = \lambda_{F,0}$ and Assumptions 1-2,*

$$GLS-LM(\lambda_{F,0}) \sim \psi' \mathcal{W}^{-1} \psi - \psi' C (C' \mathcal{W} C)^{-1} C' \psi \quad (14)$$

$$\frac{T - N + 1}{(T - K - 1)(N - K - 1)} \times JGLS(\lambda_{F,0}) \sim F_{N-K-1, T-N+1} \quad (15)$$

where $\psi \sim N(0, I_{N-1})$, $\mathcal{W} \sim \frac{\mathcal{W}_{N-1}(T-K-1, I_{N-1})}{T-K-1}$ (scaled Wishart distribution) independent of ψ , and C is an arbitrary $(N-1) \times (N-K-1)$ matrix of full rank.

Proof: See the Appendix.

The finite-sample distribution of the GLS-LM statistic is non-standard, but, since it is invariant to the specification of C , it does not depend on any further nuisance parameters so its critical values can easily be obtained by simulation.⁵ The F -distribution of the JGLS

⁵To get the finite-sample critical values for GLS-LM, we can independently simulate ψ , \mathcal{W} from normal and Wishart distributions, respectively, and use an $(N-1) \times (N-K-1)$ full rank matrix for C . The quantiles resulting from the simulated values of the GLS-LM statistic provide the critical values.

statistic has degrees of freedom different from those for the finite-sample FAR statistic, although both statistics take the limited T vs. large N problem into account.

3.2.3 FM-LM and JFM statistics

Next, we decompose the FAR statistic in an alternative manner, which is similar to the above decomposition resulting in the GLS-LM and JGLS statistics. This decomposition leads to yet two other test statistics, denoted by FM-LM and JFM respectively in Kleibergen (2009). As shown later, it yields a test whose finite-sample distribution does not depend on N .

In particular, we project $\widehat{\Sigma}^{-\frac{1}{2}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right)$ on the column space spanned by $\widehat{\Sigma}^{\frac{1}{2}}\tilde{\mathcal{B}}$ and the associated orthogonal space, to construct the FM-LM and JFM statistics, respectively. These two statistics can thus be written as:

$$\begin{aligned}
& FM-LM(\lambda_{F,0}) \\
&= \frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right)' \widehat{\Sigma}^{-\frac{1}{2}} P_{\widehat{\Sigma}^{\frac{1}{2}}\tilde{\mathcal{B}}} \widehat{\Sigma}^{-\frac{1}{2}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right) \\
&= \frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right)' \tilde{\mathcal{B}} \left(\tilde{\mathcal{B}}' \widehat{\Sigma} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right) \tag{16}
\end{aligned}$$

$$\begin{aligned}
& JFM(\lambda_{F,0}) \\
&= FAR(\lambda_{F,0}) - FM-LM(\lambda_{F,0}) \\
&= \frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right)' \widehat{\Sigma}^{-\frac{1}{2}} (I_{N-1} - P_{\widehat{\Sigma}^{\frac{1}{2}}\tilde{\mathcal{B}}}) \widehat{\Sigma}^{-\frac{1}{2}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right) \tag{17}
\end{aligned}$$

with $P_{\widehat{\Sigma}^{\frac{1}{2}}\tilde{\mathcal{B}}} = \widehat{\Sigma}^{\frac{1}{2}} \tilde{\mathcal{B}} \left(\tilde{\mathcal{B}}' \widehat{\Sigma} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \widehat{\Sigma}^{\frac{1}{2}}$, $\widehat{\Sigma} = \widehat{\Sigma}^{\frac{1}{2}} \widehat{\Sigma}^{\frac{1}{2}}$.

Theorems 5 and 6 provide the asymptotic and finite-sample distributions for these statistics, respectively.

Theorem 5. *Under $H_0 : \lambda_F = \lambda_{F,0}$ and Assumptions 1-2, the FM-LM and JFM statistics converge to independent χ_K^2 and χ_{N-K-1}^2 distributed random variables, respectively, as $T \rightarrow \infty$ while N is fixed.*

Proof: See the Appendix.

Theorem 6. Under $H_0 : \lambda_F = \lambda_{F,0}$ and Assumptions 1-2,

$$\frac{T - 2K}{(T - K - 1)K} \times FM-LM(\lambda_{F,0}) \sim F_{K, T-2K} \quad (18)$$

$$JFM(\lambda_{F,0}) \sim \psi' \mathcal{W}^{-1} \psi - \psi' \mathbf{C} (\mathbf{C}' \mathcal{W} \mathbf{C})^{-1} \mathbf{C}' \psi \quad (19)$$

where ψ , \mathcal{W} are as in Theorem 4, while \mathbf{C} is an arbitrary $(N - 1) \times K$ matrix of full rank.

Proof: See the Appendix.

It is worth noting that the finite-sample F -distribution of the FM-LM statistic in Theorem 6 has a nice feature: its degrees of freedom do not depend on the number of assets N . This statistic can thus be used to overcome the deficiency due to the presence of many assets, for which we provide simulation evidence later on.

3.2.4 Split-sample CLR (sCLR) statistic

The conditional likelihood ratio (CLR) test for $H_0 : \lambda_F = \lambda_{F,0}$ resembles the CLR test in Moreira (2003) for linear instrumental variables regression. In the single factor case, the CLR statistic has a closed form expression:

$$CLR(\lambda_{F,0}) = \frac{1}{2} \left\{ FAR(\lambda_{F,0}) - r(\lambda_{F,0}) + \sqrt{(FAR(\lambda_{F,0}) + r(\lambda_{F,0}))^2 - 4r(\lambda_{F,0})JGLS(\lambda_{F,0})} \right\} \quad (20)$$

where $r(\lambda_{F,0}) = \widehat{\mathcal{Q}}_{FF}(\lambda_{F,0}) \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}}$ corresponds to a rank test statistic for testing $\text{rank}(\mathcal{B}) = 0$ provided $\lambda_F = \lambda_{F,0}$, and it measures the strength of the risk premia identification. When $r(\lambda_{F,0})$ equals zero, so there is no identification, the CLR statistic is identical to the FAR statistic. On the other hand, for large values of $r(\lambda_{F,0})$, the CLR statistic is close to the GLS-LM statistic. Therefore, the CLR statistic can also be regarded as a data-dependent weighted average of the FAR and GLS-LM statistics.

Asymptotically, $r(\lambda_{F,0})$ is independent of $FAR(\lambda_{F,0})$ and $JGLS(\lambda_{F,0})$, so the conditional limiting distribution of the CLR statistic given the realized value of $r(\lambda_{F,0})$ results from

combining the independent limiting distributions of the GLS-LM and JGLS statistics whose sum makes up the FAR statistic. The conditional critical values of the CLR statistic can then straightforwardly be simulated for any given value of $r(\lambda_{F,0})$ in the manner of Moreira (2003), see also Kleibergen (2009). In the finite-sample setting, however, $FAR(\lambda_{F,0})$ and $JGLS(\lambda_{F,0})$ are no longer independently distributed of $r(\lambda_{F,0})$, which complicates the derivation of conditional finite-sample critical values for the CLR statistic. To address this issue, we propose to conduct a CLR test by usage of the split-sample CLR (sCLR) statistic.

Specifically, the full sample with $t = 1, \dots, T$, is divided into two parts: $t = 1, \dots, T_1$ for the first sub-sample, while $t = T_1 + 1, \dots, T$ for the second. We use the first sub-sample to estimate Σ , which is only used for calculating $r(\lambda_{F,0})$. All the other parts of the CLR statistic are computed using the second sub-sample. The resulting CLR statistic is labeled: the sCLR statistic. The $r(\lambda_{F,0})$ statistic used for the sCLR statistic is then independent of the $FAR(\lambda_{F,0})$ and $JGLS(\lambda_{F,0})$ statistics. We can thus provide the finite-sample distribution of the sCLR statistic, as in the next theorem.

Theorem 7. *Under $H_0 : \lambda_F = \lambda_{F,0}$ and Assumptions 1-2 with $K = 1$, the $sCLR(\lambda_{F,0})$ statistic conditional on the rank statistic r has the distribution:*

$$\frac{1}{2} \left(\psi' \mathcal{W}^{-1} \psi - r + \sqrt{(\psi' \mathcal{W}^{-1} \psi + r)^2 - 4r \psi' C (C' \mathcal{W} C)^{-1} C' \psi} \right)$$

where ψ , \mathcal{W} , C are as in Theorem 4, except that $T - T_1$ replaces T for the (second) sample size, and r is the rank statistic whose $\hat{\Sigma}$ results from the first sample with $t = 1, \dots, T_1$.

Proof: See the Appendix.

The split-sample CLR test is, however, not always feasible. If T is too close to N , the two split-sample sizes, T_1 and $T - T_1$, may not exceed N . Yet we require the time-series dimension to at least exceed the cross-section dimension for our proposed tests. For example, the degrees of freedom in Theorem 2 implicitly implies that we need $T - K - N + 1 > 0$. In the single factor model with $K = 1$, $T - K - N + 1 > 0$ reduces to $T > N$. For the split-sample CLR statistic, both T_1 and $T - T_1$ thus have to exceed N so T has to exceed

2*N*. For this reason, we focus on the FAR test and its components, the GLS-LM, JGLS, FM-LM and JFM tests, rather than the CLR test.

3.3 Subset FAR (sFAR) statistic

So far we have focused on testing a joint hypothesis specified on all risk premia: $H_0 : \lambda_F = \lambda_{F,0}$. To construct a confidence set of each individual risk premium in multi-factor models, we can just project out the joint confidence set which can be computed using tests of the joint hypothesis specified on all risk premia. Alternatively, Kleibergen (2009) and Kleibergen and Zhan (2019) use a subset FAR (sFAR) statistic. We provide the asymptotic and finite-sample behaviors of the sFAR statistic in Theorems 9 and 10, respectively. We refrain from constructing the asymptotic and finite-sample behaviors of the other statistics since these, unlike for the sFAR statistic, need the risk premia left unrestricted by the hypothesis of interest to be well identified.

For the $K \times 1$ dimensional vector of risk premia $\lambda_F = (\lambda_{F,1}, \lambda_{F,2}, \dots, \lambda_{F,K})'$, consider its partition such that $\lambda_F = (\lambda_{F,1}, \lambda_2)'$ with $\lambda_2 = (\lambda_{F,2}, \dots, \lambda_{F,K})'$. Our objective is to test the null hypothesis $H_0 : \lambda_{F,1} = \lambda_{F,1}^0$. The corresponding sFAR statistic, denoted by $sFAR(\lambda_{F,1}^0)$, results from minimizing the FAR statistic under the hypothesized $\lambda_{F,1}^0$:

$$sFAR(\lambda_{F,1}^0) = \min_{\lambda_2} FAR \left((\lambda_{F,1}^0, \lambda_2)' \right),$$

which equals the minimum eigenvalue of a characteristic polynomial.⁶

Proposition 8. *Suppose FAR achieves minimum at one of its infection points, then*

$$sFAR(\lambda_{F,1}^0) \leq \widetilde{JGLS} \left((\lambda_{F,1}^0, \lambda_2^0)' \right) \quad (21)$$

⁶Consider $\left| \mu \begin{pmatrix} 1 + \lambda_{F,1}' \widehat{\mathcal{Q}}_{FF,11}^{-1} \lambda_{F,1} & -\lambda_{F,1}' \widehat{\mathcal{Q}}_{FF,12}^{-1} \\ -\widehat{\mathcal{Q}}_{FF,21}^{-1} \lambda_{F,1} & \widehat{\mathcal{Q}}_{FF,22}^{-1} \end{pmatrix} - (\bar{\mathcal{R}} - \widehat{\mathcal{B}}_1 \lambda_{F,1}, \widehat{\mathcal{B}}_2)' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}} - \widehat{\mathcal{B}}_1 \lambda_{F,1}, \widehat{\mathcal{B}}_2) \right| = 0$ with $\widehat{\mathcal{Q}}_{FF}^{-1} = \begin{pmatrix} \widehat{\mathcal{Q}}_{FF,11}^{-1} & \widehat{\mathcal{Q}}_{FF,12}^{-1} \\ \widehat{\mathcal{Q}}_{FF,21}^{-1} & \widehat{\mathcal{Q}}_{FF,22}^{-1} \end{pmatrix}$, $\widehat{\mathcal{B}} = (\widehat{\mathcal{B}}_1, \widehat{\mathcal{B}}_2)$. The $sFAR(\lambda_{F,1}^0)$ statistic equals T times the smallest root. This shows the $sFAR(\lambda_{F,1}^0)$ statistic is also a rank test testing the null hypothesis that the rank of $(\mathbb{E}\mathcal{R}_t - \mathcal{B}_1 \lambda_{F,1}, \mathcal{B}_2)$ is smaller than $K_{F,2}$.

with $\widetilde{JGLS}((\lambda_{F,1}^0, \lambda_2^{0'})') = FAR((\lambda_{F,1}^0, \lambda_2^{0'})') - \widetilde{GLS - LM}((\lambda_{F,1}^0, \lambda_2^{0'})')$ where $\widetilde{GLS - LM}((\lambda_{F,1}^0, \lambda_2^{0'})')$ only uses the score with respect to the λ_2 .

Proof: See the Appendix 5.

Theorem 9. Under $H_0 : \lambda_{F,1} = \lambda_{F,1}^0$ and Assumptions 1-2, as $T \rightarrow \infty$ while N is fixed, the limiting distribution of the $sFAR(\lambda_{F,1}^0)$ statistic is bounded by the χ_{N-K}^2 distribution, i.e.,

$$sFAR(\lambda_{F,1}^0) \preceq \chi_{N-K}^2. \quad (22)$$

Proof: See the Appendix.

Theorem 10. Under $H_0 : \lambda_{F,1} = \lambda_{F,1}^0$, Assumptions 1-2, and FAR achieves minimum at one of its infection points, the scaled $sFAR(\lambda_{F,1}^0)$ statistic, $\frac{T-N}{(T-K-1)(N-K)}sFAR(\lambda_{F,1}^0)$, is in finite samples bounded by the $F_{N-K, T-N}$ distribution, i.e.,

$$\frac{T-N}{(T-K-1)(N-K)} \times sFAR(\lambda_{F,1}^0) \preceq F_{N-K, T-N}. \quad (23)$$

Proof: See the Appendix.

Theorem 11. Under $H_0 : \lambda_{F,1} = \lambda_{F,1}^0$, Assumptions 1-2, and rk_F achieves minimum at one of its infection points, then the canonical correlation rank test which tests for a lower rank value of B , rk , is in finite samples bounded by the $F_{N-K, T-N}$ distribution, i.e.,

$$\frac{T-N}{(T-K-1)(N-K)} rk \times \preceq F_{N-K, T-N}. \quad (24)$$

Proof: See the Appendix. 5

4 Numerical and empirical results

4.1 Simulation evidence

To illustrate the performance of our proposed tests, we conduct a simple simulation study. For the data generating process (DGP), we consider a single factor model:

$$R_t = c + \beta F_t + u_t, \quad \text{with } F_t \sim NID(0, V_f), \quad u_t \sim NID(0, \Omega),$$

where c , β , V_f and Ω are calibrated to data from Kroencke (2017). We set $c = \iota_N \lambda_0 + \beta \lambda_F$, so the moment condition (1) holds.

4.1.1 Size

Using the simulated data of F_t and R_t from the DGP, we test $H_0 : \lambda_F = 2$ at the 5% significance level for generating the sizes presented in Figure 1. In particular, we fix the time-series sample size at $T = 55$ to mimic the Kroencke (2017) study, while we consider a sequence of values for the number of test assets N , up to $N = 31$. The FAR, GLS-LM, JGLS, FM-LM and JFM tests described in the last section are implemented, using both their finite-sample and asymptotic critical values. The simulated sizes result from 1000 Monte Carlo replications.

Figure 1 shows that all finite-sample robust tests (dashed curves) have actual sizes close to the nominal 5%, regardless of the value of N . This is in sharp contrast with the asymptotically valid tests (solid curves), whose sizes tend to exceed the nominal 5% as N increases. When N is small, so it is far below T , we observe in Figure 1 that asymptotic tests are close to be size-correct. On the other hand, when N is large, so the limited T vs. large N problem becomes severe, asymptotic tests (except FM-LM) exhibit size distortion while finite-sample robust tests remain trustworthy. All these findings are in line with our earlier analysis.

It is worth emphasizing that the FM-LM test performs well using either finite-sample (dashed red) or asymptotic (solid red) critical values. This is explained by Theorem 6,




Figure 1: Simulated sizes as a function of the number of test assets N with $T = 55$. Dashed curves for finite-sample robust tests: FAR, dashed black; GLS-LM, dashed blue; JGLS, dashed magenta; FM-LM, dashed red; JFM, dashed green. Solid curves for asymptotically valid tests: FAR, solid black; GLS-LM, solid blue; JGLS, solid magenta; FM-LM, solid red; JFM, solid green. The null hypothesis is $H_0 : \lambda_F = 2$ in a single factor CAPM calibrated to Kroencke (2017). The significance level is 5%.




Figure 2: This figure is generated in the same manner as Figure 1 above, except that we fix T at $T = 500$.

where N does not appear in the degrees of freedom of the finite-sample distribution of the FM-LM statistic. Put differently, while the limited T vs. large N problem affects many asymptotically valid tests, it is less so for the FM-LM test. This feature makes the FM-LM test more appealing than the FAR, GLS-LM, JGLS and JFM tests for limited T vs. large N .

Figure 1 is to be compared with Figure 2, for which we increase T from 55 to 500. Under the large $T = 500$, Figure 2 shows that both the asymptotically valid tests (solid) and finite-sample robust tests (dashed) have actual sizes near the nominal 5%. This further demonstrates the importance of the T vs. N issue. If researchers adopt a large number of test assets, such as $N = 31$ assets in Kroencke (2017), or $N = 35$ as in Savov (2011), then a much larger T is generally needed to validate asymptotic tests. This argument also applies to the identification robust FAR, GLS-LM, JGLS and JFM tests when using asymptotic critical values. Kleibergen and Zhan (2019) provide similar simulation evidence for the FM t -test, whose size not only suffers from the limited T vs. large N problem, but also relies on risk factors of satisfactory quality.

4.1.2 Power

Next, we briefly analyze the power performance of our proposed tests. To do so, we vary λ_F imposed in the DGP described above and keep testing $H_0 : \lambda_F = 2$, which is close to the point estimate of the consumption risk premium reported in Kroencke (2017). To facilitate comparison, we consider two scenarios: (i) $N = 31$ and $T = 55$, as in Kroencke (2017); (ii) $N = 31$ and $T = 500$. The simulated power curves for these two scenarios are presented in Figures 3 and 4, respectively.

The left hand side panel of Figure 3 shows that finite-sample robust tests are all size-correct in our simulation study calibrated to Kroencke (2017), and these tests have good power. The power curves of the GLS-LM and FM-LM tests show a sudden decline in power which results since they are (functions of) quadratic forms of the derivative of the FAR statistic with respect to the risk premium. The power decline therefore occurs from this

derivative not only being zero at the minimizer of the FAR statistic. Andrews, Moreira and Stock (2006) show that the CLR statistic is optimal for testing in the homoskedastic linear instrumental variables regression model with one structural parameter, since it optimally combines the GLS-LM and JGLS statistic using the conditioning statistic. Since the conditioning statistic is not independently distributed from the GLS-LM and JGLS statistics, we can not use it here.

In contrast, the right hand side panel of Figure 3 shows that among the examined asymptotically valid tests, only the FM-LM test appears to be size-correct for the $N = 31$ and $T = 55$ setting. This is consistent with Figure 1 as well as Theorem 6.

Figure 3 is to be compared with Figure 4, for which we consider $T = 500$. The two panels of Figure 4 now appear similar to each other, since both the finite-sample robust tests and the asymptotic tests are expected to be valid when T is sufficiently large. Similar to the left hand side panel of Figure 3, Figure 4 also suggests that the power performances of the examined tests are comparable to each other.

Kleibergen (2009) discusses how the power decline of GLS-LM and FM-LM tests, as shown in Figure 3, can be avoided by sequentially testing the risk premia using either the independent GLS-LM and JGLS tests or the FM-LM and JFM tests. For illustration, here we consider a combination such that a 96% critical value is applied to GLS-LM and a 99% critical value to JGLS, so the overall size equals 5%. Similarly, we also consider the combined test using FM-LM (96%) and JFM (99%).

Figure 5 presents the power curves of the GLS-LM and JGLS combined test and the FM-LM and JFM combined test in the $N = 31$ and $T = 55$ setting. For comparison, we also present the power curves of the FAR test and the FM t -test with the Shanken (1992) correction.

The left hand side panel of Figure 5 shows that the power decline in Figure 3 is no longer in place, once we combine tests in the manner suggested by Kleibergen (2009). Moreover, the GLS-LM and JGLS combined test (dashed blue) appears more powerful than the FM-LM and JFM combined test (dashed red) and the FAR test (dashed black). On the other hand,




Figure 3: Simulated power curves with $N = 31$ and $T = 55$. Dashed curves for finite-sample robust tests in the left panel: FAR, dashed black; GLS-LM, dashed blue; JGLS, dashed magenta; FM-LM, dashed red; JFM, dashed green. Solid curves for asymptotically valid tests in the right panel: FAR, solid black; GLS-LM, solid blue; JGLS, solid magenta; FM-LM, solid red; JFM, solid green. The null hypothesis is $H_0 : \lambda_F = 2$ in a single factor CAPM calibrated to Kroencke (2017). The significance level is 5%.




Figure 4: This figure is generated in the same manner as Figure 3 above, except that we fix T at $T = 500$.




Figure 5: Simulated power curves with $N = 31$ and $T = 55$. Dashed curves for finite-sample robust tests in the left panel: FAR, dashed black; a combined test of GLS-LM (96%) and JGLS (99%), dashed blue; a combined test of FM-LM (96%) and JFM (99%), dashed red. Solid curves for asymptotically valid tests in the right panel: FAR, solid black; a combination of JGLS and GLS-LM, solid blue; a combination of JFM and FM-LM, solid red. In both panels, the dotted brown power curve is for the FM t -test with the Shanken (1992) correction, using critical values from the normal distribution. The null hypothesis is $H_0 : \lambda_F = 2$ in a single factor CAPM calibrated to Kroencke (2017). The significance level is 5%.




Figure 6: This figure is generated in the same manner as Figure 5 above, except that we fix T at $T = 500$.

the FM t -test (dotted brown) exhibits size distortion, since it over-rejects $H_0 : \lambda_F = 2$. The right hand side panel of Figure 5 again shows that the asymptotic counterparts of these tests do not function well in the $N = 31$ and $T = 55$ setting.

Figure 5 is to be compared with Figure 6, where T is increased to 500. Under this large value of T , the two panels in Figure 6 are, as expected, almost indistinguishable. Figure 6 also shows that the FM t -test starts to perform well under the large $T = 500$. However, its power curve lies below those of robust tests, i.e., it appears to be less powerful.

4.2 Risk premia in the consumption CAPM

We use our proposed tests to analyze risk premia in the consumption CAPM and conditional consumption CAPM. The data sets from Kroencke (2017) and Lettau and Ludvigson (2001) are adopted to facilitate comparison. The yearly data taken from Kroencke (2017) has $T = 55$ and $N = 31$, while the quarterly data taken from Lettau and Ludvigson (2001) has $T = 141$ and $N = 25$. The detailed description of the involved test assets and risk factors is provided for Tables 1-2, where the estimated β 's are reported.

For ease of exposition, we present the p -value plots resulting from testing risk premia using our proposed tests. A p -value larger than, say, 5%, implies that we could not reject the null at the 5% significance level. Thus if we find that the p -value curves are all above the 5% line, then we can not reject any hypothesized value of the risk premium, i.e., the 95% confidence sets of the risk premium is unbounded. In contrast, if the p -value curves cross the 5% line, then we can reject those hypothesized values of the risk premium whose corresponding p -values are below 5%. Finally, if the p -value curves lie fully below the 5% line, then we reject every hypothesized value of the risk premium.

Figure 7 provides the p -value curves resulting from testing the consumption risk premium using the data from Kroencke (2017). We use five of our proposed tests: FAR, GLS-LM, JGLS, FM-LM and JFM.⁷ Overall, the p -value curves based on the finite-sample robust tests (dashed black) imply unbounded 95% confidence sets of the consumption risk premium. In

⁷Since $T = 55$ while $N = 31$, we do not employ the sCLR test which needs $T > 2N$.

particular, the p -values from the finite-sample FAR, GLS-LM, JGLS and JFM tests are all above 5%, so the resulting confidence sets are unbounded. Only the FM-LM test has some of its p -values below 5%, for some hypothesized risk premia near zero. Therefore, only these hypothesized values can be rejected at the 5% level, and the 95% confidence set resulting from the FM-LM test is thus unbounded but disjoint. The unboundedness of the 95% confidence sets reflects both the questionable quality of the consumption growth data and the limited sample size, and casts doubt on the findings in Kroencke (2017) based on the FM t -test, which we re-produce in Panel A of Table 3. Dufour (1997) and Kleibergen and Zhan (2019) provide further detailed explanations for why confidence sets resulting from robust tests can be unbounded. In particular, Kleibergen and Zhan (2019) show that unbounded confidence sets commonly occur, if risk premia are unidentified due to poor quality risk factors, and/or the involved time series is short, so little information on risk premia can be extracted from the data.

Table 3: Conventional 95% confidence intervals of risk premia

	A. Kroencke (2017)		B. Lettau and Ludvigson (2001)	
	Δc	Δc	cay	$\Delta c \times \mathbf{cay}$
Estimate of λ_F	2.04	0.02	-0.13	0.06
FM t	2.18	0.20	-0.43	3.12
95% C.I. by FM t	(0.20, 3.88)	(-0.20, 0.25)	(-0.70, 0.45)	(0.02, 0.09)
Shanken t	1.75	0.15	-0.31	2.25
95% C.I. by Shanken t	(-0.24, 4.32)	(-0.29, 0.34)	(-0.93, 0.68)	(0.01, 0.11)

Notes: The estimate of λ_F and the FM t -statistic result from the Fama-MacBeth (1973) two-pass procedure. The t -statistic with the Shanken (1992) correction is also presented. 95% C.I.s are constructed by inverting the t -tests. For the consumption CAPM, we replicate the Kroencke (2017) study: the yearly unfiltered consumption growth in 1960-2014 is used for Δc , while thirty portfolios sorted by size, value and investment, plus the market portfolio, are used as test assets. For the conditional consumption CAPM, we replicate the Lettau and Ludvigson (2001) study: the quarterly consumption growth Δc , the (lagged) consumption-wealth ratio **cay** and their interaction in 1963Q3-1998Q3 are used as three risk factors, while twenty-five Fama-French portfolios sorted by size and book-to-market are used as test assets.

Figure 7 also shows the large discrepancy between finite-sample robust tests (dashed black) and asymptotically valid tests (solid red). Most striking is that the asymptotic




Figure 7: p -values of the FAR, GLS-LM, JGLS, FM-LM and JFM tests for testing the values of the consumption risk premium on the horizontal axis. Dashed black curves result from the finite-sample critical values of these test statistics, while solid red curves of p -values result from the asymptotic critical values. The test assets (thirty portfolios sorted by size, value and investment, plus the market portfolio) and the single risk factor (unfiltered consumption growth) in 1960–2014 are taken from Kroencke (2017). The 5% benchmark line is also plotted.

FAR, JGLS and JFM tests reject every hypothesized risk premium, while their finite-sample counterparts do not. The FM-LM test distinguishes itself from the other tests, since its finite-sample and asymptotic versions lead to almost identical p -values. This should not be surprising, since Theorem 6 shows that the FM-LM test does not suffer from the limited T vs. large N problem. As long as T exceeds $2K$ by a large enough margin, the finite-sample adjusted FM-LM test is almost equivalent to its asymptotic counterpart.

Identical to Figure 7, Figure 8 provides p -value plots based on the sFAR test for each of the three risk premia in the conditional consumption CAPM in Lettau and Ludvigson (2001). They are used to construct the 95% confidence sets of each of the risk premia, to contrast with the conventional ones presented in Panel B of Table 3.⁸

Figure 8 shows that the finite-sample adjusted sFAR test (dashed black) leads to unbounded 95% confidence sets of the risk premium on each of the three risk factors in the model (Δc : consumption growth; cay : the lagged consumption-wealth ratio; $\Delta c \times cay$: their interaction). In contrast, 95% confidence sets based on the asymptotic sFAR test (solid red) are unbounded and disjoint for Δc and cay , since only a bounded set of hypothesized risk premia are associated with p -values below 5%. Furthermore, for the interaction term $\Delta c \times cay$, both the finite-sample and the asymptotic p -value curves lie above 5%, so they suggest an unbounded 95% confidence set. Lettau and Ludvigson (2001), on the other hand, report a significant FM t -statistic on the interaction term to help establish their conditional consumption CAPM, which we re-produce in Panel B of Table 3. In contrast with Lettau and Ludvigson (2001), the unbounded confidence sets implied by Figure 8 thus put the significant risk premium by the FM t -test under doubt, see also Kleibergen (2009) and Kleibergen and Zhan (2019).

Finally, it is worth emphasizing that the unbounded confidence sets of risk premia in Figures 7-8 are in line with the corresponding β estimates in Tables 1-2. As indicated by Tables 1-2, we can not rule out that the full rank condition of (ι_N, β) is violated in a statistical

⁸For our other proposed tests, the 95% confidence sets for the individual risk premium result from projecting the 95% joint confidence sets of all three risk premia on the separate axes reflecting each risk premium. We therefore refrain from using them to construct the 95% confidence sets for the individual risk premium.




Figure 8: p -values of the sFAR test for testing the values of risk premia on the horizontal axis. Dashed black curves result from the sFAR test using the finite-sample F -critical values, while solid red curves of p -values result from the asymptotic χ^2 -critical values. The test assets (twenty-five Fama-French portfolios sorted by size and book-to-market ratio) and three risk factors (Δc , cay , $\Delta c \times cay$) in 1963Q3-1998Q3 are taken from Lettau and Ludvigson (2001). The 5% benchmark line is also plotted.

sense. If the full rank condition does not hold, then the risk premia λ_F in (1) are unidentified. Any real value can thus become possible for the unidentified λ_F , which consequently leads to the unbounded confidence sets in Figures 7-8.

5 Conclusions

We provide finite-sample distributions for the identification robust statistics testing risk premia in the beta representation of expected asset returns. By doing so, we make these tests suitable for empirical settings where sample sizes are often limited and the quality of risk factors is questionable. The size and power performances in Monte Carlo simulations show that these tests work favorably, and their empirical usage is illustrated using two well-known applications.

References

- [1] TW Anderson and H. Rubin. Estimation of the parameters of a single equation in a complete system of stochastic equations. *Annals of Mathematical Statistics*, pages 46–63, 1949.
- [2] Donald WK Andrews, Marcelo J Moreira, and James H Stock. Optimal two-sided invariant similar tests for instrumental variables regression. *Econometrica*, 74(3):715–752, 2006.
- [3] Marie-Claude Beaulieu, Jean-Marie Dufour, and Lynda Khalaf. Identification-robust estimation and testing of the zero-beta CAPM. *Review of Economic Studies*, 80(3):892–924, 2013.
- [4] P. Bekker. Alternative approximations to the distributions of instrumental variable estimators. *Econometrica*, pages 657–681, 1994.
- [5] P. Bekker and F. Kleibergen. Finite-sample instrumental variables inference using an asymptotically pivotal statistic. *Econometric Theory*, 19(05):744–753, 2003.
- [6] J.H. Cochrane. *Asset pricing*. Princeton University Press, 2001.
- [7] Jean-Marie Dufour. Some impossibility theorems in econometrics with applications to structural and dynamic models. *Econometrica*, pages 1365–1387, 1997.
- [8] E.F. Fama and K. French. The cross-section of expected stock returns. *Journal of Finance*, pages 427–465, 1992.
- [9] E.F. Fama and J.D. MacBeth. Risk, return, and equilibrium: Empirical tests. *Journal of Political Economy*, 81(3):607–636, 1973.
- [10] Michael R Gibbons, Stephen A Ross, and Jay Shanken. A test of the efficiency of a given portfolio. *Econometrica*, 57(5):1121–1152, 1989.
- [11] Nikolay Gospodinov, Raymond Kan, and Cesare Robotti. Spurious inference in reduced-rank asset-pricing models. *Econometrica*, 85(5):1613–1628, 2017.

- [12] Patrik Guggenberger, Frank Kleibergen, Sophocles Mavroeidis, and Linchun Chen. On the asymptotic sizes of subset anderson–rubin and lagrange multiplier tests in linear instrumental variables regression. *Econometrica*, 80(6):2649–2666, 2012.
- [13] Ravi Jagannathan and Yong Wang. Lazy investors, discretionary consumption, and the cross-section of stock returns. *Journal of Finance*, 62(4):1623–1661, 2007.
- [14] R. Kan, C. Robotti, and J. Shanken. Pricing model performance and the two-pass cross-sectional regression methodology. *Journal of Finance*, 68(6):2617–2649, 2013.
- [15] R. Kan and C. Zhang. Two-pass tests of asset pricing models with useless factors. *Journal of Finance*, pages 203–235, 1999.
- [16] R. Kan and G. Zhou. Hansen-jagannathan distance: Geometry and exact distribution. 2004.
- [17] Lynda Khalaf and Huntley Schaller. Identification and inference in two-pass asset pricing models. *Journal of Economic Dynamics and Control*, 70:165–177, 2016.
- [18] F. Kleibergen. Pivotal statistics for testing structural parameters in instrumental variables regression. *Econometrica*, pages 1781–1803, 2002.
- [19] F. Kleibergen. Tests of risk premia in linear factor models. *Journal of Econometrics*, 149(2):149–173, 2009.
- [20] F. Kleibergen and R. Paap. Generalized reduced rank tests using the singular value decomposition. *Journal of Econometrics*, 133(1):97–126, 2006.
- [21] F. Kleibergen and Z. Zhan. Unexplained factors and their effects on second pass R-squared’s. *Journal of Econometrics*, 189:101–116, 2015.
- [22] F. Kleibergen and Z. Zhan. Identification-robust inference on risk premia of mimicking portfolios of non-traded factors. *Journal of Financial Econometrics*, 16(2):155–190, 2018.

- [23] F. Kleibergen and Z. Zhan. Robust inference for consumption-based asset pricing. 2019.
- [24] Tim A Kroencke. Asset pricing without garbage. *Journal of Finance*, 72(1):47–98, 2017.
- [25] M. Lettau and S. Ludvigson. Resurrecting the (C) CAPM: A cross-sectional test when risk premia are time-varying. *Journal of Political Economy*, 109(6):1238–1287, 2001.
- [26] J. Lewellen, S. Nagel, and J. Shanken. A skeptical appraisal of asset pricing tests. *Journal of Financial Economics*, 96(2):175–194, 2010.
- [27] Thomas Mathew and Kenneth Nordström. Wishart and chi-square distributions associated with matrix quadratic forms. *Journal of Multivariate Analysis*, 61(1):129–143, 1997.
- [28] M.J. Moreira. A conditional likelihood ratio test for structural models. *Econometrica*, pages 1027–1048, 2003.
- [29] Robb J Muirhead. *Aspects of multivariate statistical theory*, volume 197. John Wiley & Sons, 2009.
- [30] W.K. Newey and F. Windmeijer. GMM with many weak moment conditions. *Econometrica*, 77(3):687–719, 2009.
- [31] Jonathan A Parker and Christian Julliard. Consumption risk and the cross section of expected returns. *Journal of Political Economy*, 113(1):185–222, 2005.
- [32] Alexi Savov. Asset pricing with garbage. *Journal of Finance*, 66(1):177–201, 2011.
- [33] J. Shanken. On the estimation of beta-pricing models. *Review of Financial Studies*, pages 1–33, 1992.

Appendix

In this Appendix, we provide the proofs for the analytical results stated in the main text. These proofs build on Lemmas 1 and 2 below.

Lemma 1. *For any given $\lambda_{F,0}$,*

$$\tilde{\mathcal{B}} = \hat{\mathcal{B}} - (\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}) \frac{\begin{pmatrix} 1 \\ -\lambda_{F,0} \end{pmatrix}' \begin{pmatrix} 1 & 0 \\ 0 & \hat{\mathcal{Q}} \end{pmatrix} \begin{pmatrix} 0 \\ I_K \end{pmatrix}}{\begin{pmatrix} 1 \\ -\lambda_{F,0} \end{pmatrix}' \begin{pmatrix} 1 & 0 \\ 0 & \hat{\mathcal{Q}} \end{pmatrix} \begin{pmatrix} 1 \\ -\lambda_{F,0} \end{pmatrix}}, \quad (25)$$

$$\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} = \frac{1}{1 + \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} \lambda_{F,0}} (\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}), \quad (26)$$

where $\hat{\mathcal{Q}} = (\sum_t \bar{F}_t \bar{F}_t') / T$.

Proof of Lemma 1:

$$\hat{\mathcal{Q}} = \left(\sum_t \bar{F}_t \bar{F}_t' \right) / T \quad (27)$$

$$\hat{\mathcal{Q}}_{FF}(\lambda_{F,0}) = \left(\sum_t (\bar{F}_t + \lambda_{F,0})(\bar{F}_t + \lambda_{F,0})' \right) / T = \hat{\mathcal{Q}} + \lambda_{F,0} \lambda'_{F,0} \quad (28)$$

where the last equality results from the fact that \bar{F}_t are demeaned factors. Using $(A + aa')^{-1} = A^{-1} - \frac{1}{1+a'a} A^{-1} aa'^{-1}$, we have

$$\hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} = \left(\hat{\mathcal{Q}} + \lambda_{F,0} \lambda'_{F,0} \right)^{-1} = \hat{\mathcal{Q}}^{-1} - \frac{1}{1 + \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} \lambda_{F,0}} \hat{\mathcal{Q}}^{-1} \lambda_{F,0} \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} \quad (29)$$

which implies

$$1 - \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} \lambda_{F,0} = \frac{1}{1 + \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} \lambda_{F,0}} \quad (30)$$

Plugging Equation (29) into $\tilde{\mathcal{B}}$, which can be rewritten as $\tilde{\mathcal{B}} = \frac{1}{T}\mathcal{R}'(\iota_T\lambda'_{F,0} + F)\hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1}$ with $F = (\bar{F}_1, \dots, \bar{F}_T)'$ and $\mathcal{R} = (\mathcal{R}_1, \dots, \mathcal{R}_T)'$, we get:

$$\begin{aligned}
\tilde{\mathcal{B}} &= \frac{1}{T}\mathcal{R}'(\iota_T\lambda'_{F,0} + F)\hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} \\
&= \frac{1}{T}\mathcal{R}'F\left(\hat{\mathcal{Q}}^{-1} - \frac{1}{1 + \lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}\lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\right) + \bar{\mathcal{R}}\lambda'_{F,0}\hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} \\
&= \hat{\mathcal{B}} - \frac{1}{1 + \lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}}\hat{\mathcal{B}}\lambda_{F,0}\lambda'_{F,0}\hat{\mathcal{Q}}^{-1} + \bar{\mathcal{R}}\lambda'_{F,0}\hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} \\
&= \hat{\mathcal{B}} - \frac{1}{1 + \lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}}\hat{\mathcal{B}}\lambda_{F,0}\lambda'_{F,0}\hat{\mathcal{Q}}^{-1} + \bar{\mathcal{R}}\lambda'_{F,0}\left(\hat{\mathcal{Q}}^{-1} - \frac{1}{1 + \lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}\lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\right) \\
&= \hat{\mathcal{B}} - \frac{1}{1 + \lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}}\hat{\mathcal{B}}\lambda_{F,0}\lambda'_{F,0}\hat{\mathcal{Q}}^{-1} + \frac{1}{1 + \lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}}\bar{\mathcal{R}}\lambda'_{F,0}\hat{\mathcal{Q}}^{-1} \\
&= \hat{\mathcal{B}} - \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}\right)\frac{-\lambda'_{F,0}\hat{\mathcal{Q}}^{-1}}{1 + \lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}}
\end{aligned} \tag{31}$$

which then implies the following Equations (32) and (33):

$$\tilde{\mathcal{B}} = \hat{\mathcal{B}} - \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}\right)\frac{\begin{pmatrix} 1 \\ -\lambda_{F,0} \end{pmatrix}' \begin{pmatrix} 1 & 0 \\ 0 & \hat{\mathcal{Q}}^{-1} \end{pmatrix} \begin{pmatrix} 0 \\ I_K \end{pmatrix}}{\begin{pmatrix} 1 \\ -\lambda_{F,0} \end{pmatrix}' \begin{pmatrix} 1 & 0 \\ 0 & \hat{\mathcal{Q}}^{-1} \end{pmatrix} \begin{pmatrix} 1 \\ -\lambda_{F,0} \end{pmatrix}} \tag{32}$$

$$\tilde{\mathcal{B}}\lambda_{F,0} = \hat{\mathcal{B}}\lambda_{F,0} - \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}\right)\frac{-\lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}}{1 + \lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}} \tag{33}$$

From Equation (33), it results that

$$\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} = \frac{1}{1 + \lambda'_{F,0}\hat{\mathcal{Q}}^{-1}\lambda_{F,0}}\left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}\right) \tag{34}$$

Q.E.D.

Remarks. Lemma 1 also implies

$$\tilde{\mathcal{B}} - \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right) \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} = \hat{\mathcal{B}}. \quad (35)$$

If the model is correctly specified, the term $\left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right)$ is asymptotically zero, so the two estimators $\tilde{\mathcal{B}}$ and $\hat{\mathcal{B}}$ are asymptotically equivalent.

Lemma 2. *Under $H_0 : \lambda_F = \lambda_{F,0}$ and Assumption1 1-2,*

$$\sqrt{T} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0} \right) \sim N \left(0, \left(1 + \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} \lambda_{F,0} \right) \otimes \Sigma \right), \quad (36)$$

$$\sqrt{T} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right) \sim N \left(0, \left(1 - \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} \lambda_{F,0} \right) \otimes \Sigma \right), \quad (37)$$

$$\left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0} \right) \perp \tilde{\mathcal{B}}, \quad (38)$$

$$\left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right) \perp \tilde{\mathcal{B}}, \quad (39)$$

$$\left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0} \right) \perp \hat{\Sigma}, \quad (40)$$

$$\left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right) \perp \hat{\Sigma}, \quad (41)$$

$$\hat{\Sigma} = \frac{1}{T - K - 1} \sum_t \left(\bar{\mathcal{R}}_t - \hat{\mathcal{B}}\bar{F}_t \right) \left(\bar{\mathcal{R}}_t - \hat{\mathcal{B}}\bar{F}_t \right)' \sim \frac{\mathcal{W}_{N-1}(T - K - 1, \Sigma)}{(T - K - 1)}, \quad (42)$$

$$\bar{\mathcal{R}} \perp \hat{\mathcal{B}}, \bar{\mathcal{R}} \perp \hat{\Sigma}, \hat{\mathcal{B}} \perp \hat{\Sigma}, \tilde{\mathcal{B}} \perp \hat{\Sigma}. \quad (43)$$

Proof of Lemma 2: We only need to show (36) (39) (40)(42). The proof of these results, together with Lemma 1, implies the rest of Lemma 2.

(1) Show $\sqrt{T} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0} \right) \sim N \left(0, \left(1 + \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} \lambda_{F,0} \right) \otimes \Sigma \right)$:

$$\begin{aligned} \sqrt{T} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0} \right) &= \sqrt{T} \left(\bar{\mathcal{R}} - \mathcal{B}\lambda_{F,0} - \left(\hat{\mathcal{B}} - \mathcal{B} \right) \lambda_{F,0} \right) \\ &= \sqrt{T} \left(\bar{\mathcal{U}} - \left(\hat{\mathcal{B}} - \mathcal{B} \right) \lambda_{F,0} \right) \\ &= \sqrt{T} \left(\mathcal{U}' \iota_T (\iota_T' \iota_T)^{-1} - \mathcal{U}' F (F' F)^{-1} \lambda_{F,0} \right) \\ &= \sqrt{T} \mathcal{U}' X (X' X)^{-1} \begin{pmatrix} 1 & -\lambda'_{F,0} \end{pmatrix}' \end{aligned} \quad (44)$$

where $\mathcal{U} = (\mathcal{U}_1, \dots, \mathcal{U}_T)'$, $F = (\bar{F}_1, \dots, \bar{F}_T)'$ and $X = (\iota_T, F)$. It then follows that

$$\begin{aligned} \sqrt{T} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_{F,0} \right) &= \sqrt{T} \mathcal{U}' X (X' X)^{-1} \begin{pmatrix} 1 & -\lambda'_{F,0} \end{pmatrix}' \\ &= \left(\begin{pmatrix} 1 & -\lambda'_{F,0} \end{pmatrix} (X' X / T)^{-1} \otimes I_{N-1} \right) \left(\frac{1}{\sqrt{T}} \sum_t X_t \otimes \mathcal{U}_t \right) \end{aligned} \quad (45)$$

and we thus get (36) by using

$$\frac{1}{\sqrt{T}} \sum_t X_t \otimes \mathcal{U}_t \sim N(0, (X' X) / T \otimes \Sigma) \quad (46)$$

(2) Show $(\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0}) \perp \tilde{\mathcal{B}}$, where \perp stands for independence:

We prove the claim by showing

$$\begin{pmatrix} \bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \\ \text{vec}(\tilde{\mathcal{B}} - \mathcal{B}) \end{pmatrix} \sim \frac{1}{\sqrt{T}} \begin{pmatrix} \psi_{\mathcal{R}} \\ \psi_{\mathcal{B}} \end{pmatrix} \quad (47)$$

where $\psi_{\mathcal{R}} \perp \psi_{\mathcal{B}}$, and they are $(N-1) \times 1$ and $(N-1)K \times 1$ normally distributed random vectors with mean zero and covariance matrices $(1 - \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}) \otimes \Sigma$ and $\hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} \otimes \Sigma$, respectively. To show it, we start with

$$\begin{pmatrix} \bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \\ \text{vec}(\tilde{\mathcal{B}} - \mathcal{B}) \end{pmatrix} \sim \begin{pmatrix} \bar{\mathcal{R}} - \mathcal{B} \lambda_{F,0} - (\lambda'_{F,0} \otimes I_{N-1}) \text{vec}(\tilde{\mathcal{B}} - \mathcal{B}) \\ \text{vec}(\tilde{\mathcal{B}} - \mathcal{B}) \end{pmatrix} \quad (48)$$

where

$$\bar{\mathcal{R}} - \mathcal{B} \lambda_{F,0} = \bar{\mathcal{U}} = \mathcal{U}' \iota_T (\iota_T' \iota_T)^{-1} = ((\iota_T' \iota_T)^{-1} \iota_T' \otimes I_{N-1}) \text{vec}(\mathcal{U}') \quad (49)$$

$$\text{vec}(\tilde{\mathcal{B}} - \mathcal{B}) = \frac{1}{T} (\hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) (F' + \lambda_{F,0} \iota_T') \otimes I_{N-1}) \text{vec}(\mathcal{U}') \quad (50)$$

We therefore have

$$\begin{aligned}
\begin{pmatrix} \bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \\ \text{vec}(\tilde{\mathcal{B}} - \mathcal{B}) \end{pmatrix} &\sim \begin{pmatrix} I_{N-1} & -(\lambda'_{F,0} \otimes I_{N-1}) \\ 0 & I_{K(N-1)} \end{pmatrix} \begin{pmatrix} (\iota'_T \iota_T)^{-1} \iota'_T \otimes I_{N-1} \\ \frac{1}{T} \hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) (F + \iota_T \lambda'_{F,0})' \otimes I_{N-1} \end{pmatrix} \text{vec}(\mathcal{U}') \\
&\sim \begin{pmatrix} \left((\iota'_T \iota_T)^{-1} \iota'_T - \frac{1}{T} \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) (F + \iota_T \lambda'_{F,0})' \right) \otimes I_{N-1} \\ \frac{1}{T} \hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) (F + \iota_T \lambda'_{F,0})' \otimes I_{N-1} \end{pmatrix} \text{vec}(\mathcal{U}') \\
&\sim \frac{1}{T} \begin{pmatrix} \left(\iota'_T - \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) (F + \iota_T \lambda'_{F,0})' \right) \otimes I_{N-1} \\ \hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) (F + \iota_T \lambda'_{F,0})' \otimes I_{N-1} \end{pmatrix} \text{vec}(\mathcal{U}') \quad (51)
\end{aligned}$$

Note that $\left(\iota'_T - \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) (F + \iota_T \lambda'_{F,0})' \right) \times \left(\hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) (F + \iota_T \lambda'_{F,0})' \right)' = 0$, which implies $\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \perp \tilde{\mathcal{B}}$.

(3) Show $(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}) \perp \hat{\Sigma}$:

Note that $\hat{\Sigma} = \mathcal{U}' M_X \mathcal{U} / (T - K - 1)$ with $M_X = I_T - X(X'X)^{-1}X'$, while $(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0}) = \mathcal{U}' X (X'X)^{-1} \begin{pmatrix} 1 & -\lambda'_{F,0} \end{pmatrix}'$ as shown in (44), the independence thus results from $\mathcal{U}' X \perp \mathcal{U}' M_X$, since

$$\mathbb{E}(\text{vec}(\mathcal{U}' X)' \text{vec}(\mathcal{U}' M_X)) = \mathbb{E}((X' \otimes I_{N-1}) \text{vec}(\mathcal{U}')' (M_X \otimes I_{N-1}) \text{vec}(\mathcal{U}' M_X)) = 0 \quad (52)$$

(4) Show $\hat{\Sigma} \sim \frac{\mathcal{W}_{N-1}(T-K-1, \Sigma)}{(T-K-1)}$:

$\hat{\Sigma} = \mathcal{U}' M_X \mathcal{U} / (T - K - 1)$, where $M_X = I_T - X(X'X)^{-1}X'$ is the projection matrix with rank $T - K - 1$, $\mathcal{U} = (\mathcal{U}_1, \dots, \mathcal{U}_T)'$ with $\mathcal{U}_t \sim N(0_{N-1}, \Sigma)$. Using Theorem 2 of Matthew and Nordstrom (1997), it follows that

$$\hat{\Sigma} = \mathcal{U}' M_X \mathcal{U} / (T - K - 1) \sim \mathcal{W}_{N-1}(T - K - 1, \Sigma) / (T - K - 1). \quad (53)$$

Q.E.D.

Proof of Theorem 1: Equivalence of FAR and H statistics results from (26) (30):

$$\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} = \frac{1}{1 + \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} \lambda_{F,0}} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0} \right) \quad (54)$$

$$1 - \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} \lambda_{F,0} = \frac{1}{1 + \lambda'_{F,0} \hat{\mathcal{Q}}^{-1} \lambda_{F,0}} \quad (55)$$

Assumptions 1 and 2 imply that Assumption 1 in Kleibergen (2009) holds. Theorem 1 thus results from Theorem 5 in Kleibergen (2009).

Q.E.D.

Proof of Theorem 2: From Lemma 2, we know:

$$\sqrt{\frac{T}{1 - \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} \lambda_{F,0}}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_{F,0} \right) \sim N(0, \Sigma) \quad (56)$$

$$\left(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_{F,0} \right) \perp \hat{\Sigma} \quad (57)$$

$$\hat{\Sigma} = \frac{1}{T - K - 1} \sum_t \left(\bar{\mathcal{R}}_t - \hat{B}\bar{F}_t \right) \left(\bar{\mathcal{R}}_t - \hat{B}\bar{F}_t \right)' \sim \frac{\mathcal{W}_{N-1}(T - K - 1, \Sigma)}{(T - K - 1)} \quad (58)$$

With these results, the $F_{N-1, T-K-N+1}$ distribution for the finite-sample FAR statistic results from Theorem 3.2.13 from Muirhead (2009).

Q.E.D.

Proof of Theorem 3: Similar to the proof of Theorem 1, Theorem 3 results from Theorem 6 in Kleibergen (2009).

Q.E.D.

Proof of Theorem 4: Let $\tilde{\mathcal{B}}_{\perp}$ be an arbitrary $(N - 1) \times (N - K - 1)$ matrix such that $(\tilde{\mathcal{B}}, \tilde{\mathcal{B}}_{\perp})$ is of full rank and $\tilde{\mathcal{B}}'_{\perp} \tilde{\mathcal{B}} = 0$. The projection matrix $M_{\hat{\Sigma}^{-\frac{1}{2}} \tilde{\mathcal{B}}} = I_{N-1} - P_{\hat{\Sigma}^{-\frac{1}{2}} \tilde{\mathcal{B}}}$ can be written as $\hat{\Sigma}^{\frac{1}{2}} \tilde{\mathcal{B}}_{\perp} \left(\tilde{\mathcal{B}}'_{\perp} \hat{\Sigma} \tilde{\mathcal{B}}_{\perp} \right)^{-1} \tilde{\mathcal{B}}'_{\perp} \hat{\Sigma}^{\frac{1}{2}}$.

The $JGLS(\lambda_{F,0})$ statistic can thus be written as:

$$\begin{aligned}
& JGLS(\lambda_{F,0}) \\
&= \frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \right)' \widehat{\Sigma}^{-\frac{1}{2}'} (I_{N-1} - P_{\widehat{\Sigma}^{-\frac{1}{2}} \tilde{\mathcal{B}}}) \widehat{\Sigma}^{-\frac{1}{2}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \right) \\
&= \frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \right)' \tilde{\mathcal{B}}_{\perp} \left(\tilde{\mathcal{B}}'_{\perp} \widehat{\Sigma} \tilde{\mathcal{B}}_{\perp} \right)^{-1} \tilde{\mathcal{B}}'_{\perp} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \right) \quad (59)
\end{aligned}$$

Conditional on $\tilde{\mathcal{B}}$, Lemma 2 implies:

$$\sqrt{\frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}}} \tilde{\mathcal{B}}'_{\perp} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \right) \sim N \left(0, \tilde{\mathcal{B}}'_{\perp} \Sigma \tilde{\mathcal{B}}_{\perp} \right) \quad (60)$$

$$\tilde{\mathcal{B}}'_{\perp} \widehat{\Sigma} \tilde{\mathcal{B}}_{\perp} \sim \frac{\mathcal{W}_{N-K-1} \left(T - K - 1, (\tilde{\mathcal{B}}_{\perp})' \Sigma (\tilde{\mathcal{B}}_{\perp}) \right)}{T - K - 1} \quad (61)$$

$$\tilde{\mathcal{B}}'_{\perp} \widehat{\Sigma} \tilde{\mathcal{B}}_{\perp} \perp \tilde{\mathcal{B}}'_{\perp} \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \right) \quad (62)$$

since $\tilde{\mathcal{B}}_{\perp}$ only depends on $\tilde{\mathcal{B}}$ by construction. These results imply that $\frac{T-N+1}{(T-K-1)(N-K-1)} \times JGLS(\lambda_{F,0}) \sim F_{N-K-1, T-N+1}$.

Note that $GLS-LM(\lambda_{F,0}) = FAR(\lambda_{F,0}) - JGLS(\lambda_{F,0})$, where $FAR(\lambda_{F,0})$ and $JGLS(\lambda_{F,0})$, once properly scaled, follow F -distributions with different degrees of freedom. The finite distribution of $GLS-LM(\lambda_{F,0})$ is thus the difference of finite-sample distributions of $FAR(\lambda_{F,0})$ and $JGLS(\lambda_{F,0})$ statistics. From the proof of Lemma 2, we know

$$\begin{aligned}
\sqrt{T} \begin{pmatrix} \bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \\ \text{vec} \left(\tilde{\mathcal{B}} - \mathcal{B} \right) \end{pmatrix} &\sim \begin{pmatrix} \psi_{\mathcal{R}} \\ \psi_{\mathcal{B}} \end{pmatrix} \\
&\sim N \left(0, \begin{pmatrix} \left(1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0} \right) \otimes \Sigma & 0 \\ 0 & \widehat{\mathcal{Q}}_{FF}(\lambda_{F,0})^{-1} \otimes \Sigma \end{pmatrix} \right) \quad (63)
\end{aligned}$$

and $\widehat{\Sigma} \sim \frac{\mathcal{W}_{N-1}(T-K-1, \Sigma)}{(T-K-1)}$, $\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0} \perp \widehat{\Sigma}$. Therefore, we have $FAR(\lambda_{F,0}) \sim \psi' \mathcal{W}^{-1} \psi$, where $\psi \sim N(0, I_{N-1})$, $\mathcal{W} \sim \frac{\mathcal{W}_{N-1}(T-K-1, I_{N-1})}{T-K-1}$. As $\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0}, \widehat{\Sigma}$ are independent from $\tilde{\mathcal{B}}_{\perp}$, we have

$JGLS(\lambda_{F,0}) \sim \psi' C (C' \mathcal{W} C)^{-1} C' \psi$ independent on C from (59).

Q.E.D.

Proof of Theorem 5: Similar to the proof of Theorem 1, Theorem 5 results from Theorem 7 in Kleibergen (2009).

Q.E.D.

Proof of Theorem 6: Conditional on $\tilde{\mathcal{B}}$, Lemma 2 implies:

$$\sqrt{\frac{T}{1 - \lambda'_{F,0} \hat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}}} \tilde{\mathcal{B}}' (\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0}) \sim N(0, \tilde{\mathcal{B}}' \Sigma \tilde{\mathcal{B}}) \quad (64)$$

$$\tilde{\mathcal{B}}' \hat{\Sigma} \tilde{\mathcal{B}} \sim \frac{\mathcal{W}_K(T - K - 1, \tilde{\mathcal{B}}' \Sigma \tilde{\mathcal{B}})}{T - K - 1} \quad (65)$$

$$\tilde{\mathcal{B}}' \hat{\Sigma} \tilde{\mathcal{B}} \perp \tilde{\mathcal{B}}' (\bar{\mathcal{R}} - \tilde{\mathcal{B}} \lambda_{F,0}) \quad (66)$$

These results imply that $\frac{T-2K}{(T-K-1)K} \times FM-LM(\lambda_{F,0}) \sim F_{K,T-2K}$, and this distribution does not depend on $\tilde{\mathcal{B}}$.

Given $JFM(\lambda_{F,0}) = FAR(\lambda_{F,0}) - FM-LM(\lambda_{F,0})$, the finite distribution of $JFM(\lambda_{F,0})$ is thus the difference of finite-sample distributions of $FAR(\lambda_{F,0})$ and $FM-LM(\lambda_{F,0})$ statistics: $JFM(\lambda_{F,0}) \sim \psi' \mathcal{W}^{-1} \psi - \psi' \mathbf{C} (\mathbf{C}' \mathcal{W} \mathbf{C})^{-1} \mathbf{C}' \psi$. The proof is similar to that for GLS-LM in Theorem 4, except for that \mathbf{C} is an arbitrary $(N - 1) \times K$ matrix of full rank.

Q.E.D.

Proof of Theorem 7: The i.i.d. condition in Assumption 1 implies that r in the sCLR statistic is independent from the other parts. Theorem 7 thus results from Lemma 2 and Theorem 4.

Q.E.D.

Some extra remarks before proof of Proposition 8:

(1)

$$GLS - LM(\lambda_F^0) = \frac{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F}{4T} \frac{\partial FAR(\lambda_F)}{\partial \lambda'_F} \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \left(\frac{\partial FAR(\lambda_F)}{\partial \lambda'_F} \right)' \quad (67)$$

(2)

$$\begin{aligned} \widetilde{GLS} - LM \left(\left(\lambda_{F,1}^0, \lambda_2^{0'} \right)' \right) &= \frac{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F}{4T} \frac{\partial FAR(\lambda_F)}{\partial \lambda'_2} \Big|_{\lambda_F = \lambda_F^0} \left(\widetilde{\mathcal{B}}_2' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}}_2 \right)^{-1} \left(\frac{\partial FAR(\lambda_F)}{\partial \lambda'_2} \Big|_{\lambda_F = \lambda_F^0} \right)' \\ &= \frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} \left(\bar{\mathcal{R}} - \widetilde{\mathcal{B}} \lambda_{F,0} \right)' \widehat{\Sigma}^{-\frac{1}{2}'} P_{\widehat{\Sigma}^{-\frac{1}{2}} \widetilde{\mathcal{B}}_2} \widehat{\Sigma}^{-\frac{1}{2}} \left(\bar{\mathcal{R}} - \widetilde{\mathcal{B}} \lambda_{F,0} \right) \end{aligned} \quad (68)$$

(3)

$$\widetilde{JGLS} \left(\left(\lambda_{F,1}^0, \hat{\lambda}'_2(\lambda_{F,1}^0) \right)' \right) = \frac{T}{1 - \lambda'_{F,0} \widehat{\mathcal{Q}}_{FF}^{-1}(\lambda_{F,0}) \lambda_{F,0}} \left(\bar{\mathcal{R}} - \widetilde{\mathcal{B}} \lambda_{F,0} \right)' \widehat{\Sigma}^{-\frac{1}{2}'} (I_{N-1} - P_{\widehat{\Sigma}^{-\frac{1}{2}} \widetilde{\mathcal{B}}_2}) \widehat{\Sigma}^{-\frac{1}{2}} \left(\bar{\mathcal{R}} - \widetilde{\mathcal{B}} \lambda_{F,0} \right) \quad (69)$$

$$\begin{aligned} \frac{\partial FAR(\lambda_F)}{\partial \lambda'_F} &= \frac{T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \frac{\partial \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)}{\partial \lambda'_F} \\ &\quad + \frac{\partial \frac{T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F}}{\partial \lambda'_F} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right) \\ &= \frac{T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} 2 \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \left(-\widehat{\mathcal{B}} \right) - \frac{T}{(1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} 2 \lambda'_F \widehat{\mathcal{Q}}^{-1} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right) \\ &= - \frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \left(\widehat{\mathcal{B}} - \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right) \frac{-\lambda'_F \widehat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \right) \\ &= - \frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} = -2T \left(\bar{\mathcal{R}} - \widetilde{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \end{aligned} \quad (70)$$

and thus

$$GLS-LM(\lambda_F) = \frac{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F}{4T} \frac{\partial FAR(\lambda_F)}{\partial \lambda'_F} \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \left(\frac{\partial FAR(\lambda_F)}{\partial \lambda'_F} \right)' \quad (71)$$

Similarly we have

$$\frac{\partial FAR(\lambda_F)}{\partial \lambda'_2} = -2T \left(\bar{\mathcal{R}} - \tilde{\mathcal{B}}\lambda_F \right)' \hat{\Sigma}^{-1} \tilde{\mathcal{B}}_2 \quad (72)$$

Proof of Proposition 8: Denote

$$\hat{\lambda}_2(\lambda_{F,1}^0) = \arg \min_{\lambda_2} FAR \left((\lambda_{F,1}^0, \lambda'_2)' \right)$$

By construction $\widetilde{GLS} - LM \left((\lambda_{F,1}^0, \hat{\lambda}'_2(\lambda_{F,1}^0))' \right) = 0$ and thus once we establish that $\widetilde{JGLS} \left((\lambda_{F,1}^0, \hat{\lambda}'_2(\lambda_{F,1}^0))' \right) \leq \widetilde{JGLS} \left((\lambda_{F,1}^0, \lambda_2^{0'})' \right)$ we have

$$sFAR(\lambda_{F,1}^0) = \widetilde{JGLS} \left((\lambda_{F,1}^0, \hat{\lambda}'_2(\lambda_{F,1}^0))' \right) \leq \widetilde{JGLS} \left((\lambda_{F,1}^0, \lambda_2^{0'})' \right)$$

The inequality $\widetilde{JGLS} \left((\lambda_{F,1}^0, \hat{\lambda}'_2(\lambda_{F,1}^0))' \right) \leq \widetilde{JGLS} \left((\lambda_{F,1}^0, \lambda_2^{0'})' \right)$ holds since the derivative of $\widetilde{JGLS} \left((\lambda_{F,1}^0, \lambda'_2)' \right)$ with respect to λ_2 hits zero at $\hat{\lambda}'_2(\lambda_{F,1}^0)$. Given the facts that stationary points of FAR and \widetilde{JGLS} coincide, they achieve the minimum at the same point.

Q.E.D.

Proof of Theorem 9: Similar to the proof of Theorem 1, Theorem 9 results from Theorem 9 in Kleibergen (2009). See also Guggenberger et al. (2012), Gospodinov et al. (2017).

Q.E.D.

Proof of Theorem 10: This is a direct result of Proposition 8.??

The sFAR statistic equals T times the smallest root of the following characteristic polynomial:

$$\left| \mu \begin{pmatrix} 1 + \lambda'_{F,1} \hat{\mathcal{Q}}_{FF,11}^{-1} \lambda_{F,1} & -\lambda'_{F,1} \hat{\mathcal{Q}}_{FF,12}^{-1} \\ -\hat{\mathcal{Q}}_{FF,21}^{-1} \lambda_{F,1} & \hat{\mathcal{Q}}_{FF,22}^{-1} \end{pmatrix} - \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}_1 \lambda_{F,1}, \hat{\mathcal{B}}_2 \right)' \hat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}}_1 \lambda_{F,1}, \hat{\mathcal{B}}_2 \right) \right| = 0$$

with $\widehat{Q}_{FF}^{-1} = \begin{pmatrix} \widehat{Q}_{FF,11}^{-1} & \widehat{Q}_{FF,12}^{-1} \\ \widehat{Q}_{FF,21}^{-1} & \widehat{Q}_{FF,22}^{-1} \end{pmatrix}$, $\widehat{B} = (\widehat{B}_1, \widehat{B}_2)$, $\lambda_{F,1}$ results from its hypothesized value, and $\lambda_F = (\lambda_{F,1}, \lambda_2)'$.

Let $\bar{\mathcal{R}}_1$ denote $\bar{\mathcal{R}} - \widehat{B}_1 \lambda_{F,1}$, so the polynomial above can be expressed as

$$\left| \mu \begin{pmatrix} 1 + \lambda'_{F,1} \widehat{Q}_{FF,11}^{-1} \lambda_{F,1} & -\lambda'_{F,1} \widehat{Q}_{FF,12}^{-1} \\ -\widehat{Q}_{FF,21}^{-1} \lambda_{F,1} & \widehat{Q}_{FF,22}^{-1} \end{pmatrix} - (\bar{\mathcal{R}}_1, \widehat{B}_2)' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}}_1, \widehat{B}_2) \right| = 0,$$

which shows that the sFAR statistic is also the minimum of the FAR statistic for testing λ_2 , given $\lambda_{F,1}$.

Recall that the FAR statistic can be decomposed to JGLS and GLS-LM statistics; in addition, the GLS-LM statistic equals the quadratic form of the derivative of the FAR statistic. This implies that the minimum of FAR coincides with the minimum of JGLS. Therefore, the sFAR statistic is upper bounded by the distribution of JGLS for testing λ_2 provided $\lambda_{F,1}$. With the proper scaling, the exact finite-sample distribution of such a JGLS statistic by Theorem 4 is as $F_{N-K, T-N}$ (note: $K - 1$ now plays the role of K in Theorem 4, since the dimension of λ_2 is $(K - 1) \times 1$). Consequently, the scaled sFAR statistic is bounded by this F -distribution.

Q.E.D.

Proof of Theorem 11: By construction we know $\widehat{B} \sim N(B, Q_F^{-1} \otimes \Sigma)$. Suppose H_0 : B is of lower rank than K then the null is equivalent to that there exists λ_0 such that

$B(1, -\lambda'_0)' = B_1 - B_2\lambda_0 = 0$. Denote

$$\begin{aligned}\tilde{\mathcal{B}}_2 &= \hat{\mathcal{B}}_2 + \hat{\mathcal{B}}(1, -\lambda'_0)' A \\ rk_F &= \frac{T}{(1, -\lambda'_0)' Q (1, -\lambda'_0)'} \left(\hat{\mathcal{B}}(1, -\lambda'_0)' \right)' \hat{\Sigma}^{-1} \left(\hat{\mathcal{B}}(1, -\lambda'_0)' \right) \\ rk_{\tilde{G}} &= \frac{T}{(1, -\lambda'_0)' Q (1, -\lambda'_0)'} \left(\hat{\mathcal{B}}(1, -\lambda'_0)' \right)' \hat{\Sigma}^{-\frac{1}{2}} P_{\hat{\Sigma}^{-\frac{1}{2}} \tilde{\mathcal{B}}_2} \hat{\Sigma}^{-\frac{1}{2}} \left(\hat{\mathcal{B}}(1, -\lambda'_0)' \right) \\ rk_{\tilde{J}} &= rk_F - rk_{\tilde{G}}\end{aligned}$$

$$\text{with } Q = Q_F^{-1}, A = \frac{\begin{pmatrix} 1 \\ -\lambda_0 \end{pmatrix}' Q \begin{pmatrix} 0 \\ I_{K-1} \end{pmatrix}}{\begin{pmatrix} 1 \\ -\lambda_0 \end{pmatrix}' Q \begin{pmatrix} 1 \\ -\lambda_0 \end{pmatrix}} ..$$

Then the result follows proof of Theorem 10 by noticing $\min_{\lambda} rk_F(\lambda) \leq rk_{\tilde{J}}(\lambda_0)$ and $\min_{\lambda} rk_F(\lambda)$ equals T times the minimal root of the following polynomial, which is the canonical correlation rank test

$$\left| \mu Q - \hat{\mathcal{B}}' \hat{\Sigma}^{-1} \hat{\mathcal{B}} \right| = 0,$$

Here we briefly illustrate the fact that $\tilde{\mathcal{B}}_2 \perp \hat{\mathcal{B}}(1, -\lambda'_0)'$.

$$\left(\hat{\mathcal{B}}(1, -\lambda'_0)', \tilde{\mathcal{B}}_2 \right) = \hat{\mathcal{B}} A_{\lambda_0} \tag{73}$$

$$\text{with } A_{\lambda_0} = \begin{pmatrix} 1 & A \\ -\lambda_0 & I_{K-1} - \lambda_0 A \end{pmatrix}. \{AA'_{\lambda_0} Q A_{\lambda_0}\}_{12} = 0 \text{ implies the result.}$$

Q.E.D.

6 Supplementary part

Proposition 3. *The GLS-LM(λ_F) statistic (12) is a quadratic form of the derivative of FAR(λ_F).*

Proof of proposition 3:

$$\begin{aligned}
\frac{\partial FAR(\lambda_F)}{\partial \lambda'_F} &= \frac{T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \frac{\partial \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)}{\partial \lambda'_F} \\
&+ \frac{\partial \frac{T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F}}{\partial \lambda'_F} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right) \\
&= \frac{T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} 2 \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \left(-\widehat{\mathcal{B}} \right) - \frac{T}{(1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} 2 \lambda'_F \widehat{\mathcal{Q}}^{-1} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right) \\
&= - \frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \left(\widehat{\mathcal{B}} - \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right) \frac{-\lambda'_F \widehat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \right) \\
&= - \frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} = - - 2T \left(\bar{\mathcal{R}} - \widetilde{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \tag{74}
\end{aligned}$$

and thus

$$\text{GLS-LM}(\lambda_F) = \frac{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F}{4T} \frac{\partial FAR(\lambda_F)}{\partial \lambda'_F} \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \left(\frac{\partial FAR(\lambda_F)}{\partial \lambda'_F} \right)' \tag{75}$$

Q.E.D

Proposition 4.

$$\frac{\partial \text{GLS-LM}}{\partial \lambda'_F}(\lambda_F) = \frac{\partial FAR}{\partial \lambda'_F} - \frac{J\text{GLS}(\lambda_F)}{T} \frac{\partial FAR}{\partial \lambda'_F} \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \mathcal{Q}_- \tag{76}$$

where $\mathcal{Q}_- = \left(\widehat{\mathcal{Q}}^{-1} - \frac{\widehat{\mathcal{Q}}^{-1} \lambda_F \lambda'_F \widehat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \right)$.

Proof of proposition 4:

$$\begin{aligned}
d\tilde{\mathcal{B}} &= d \left(\hat{\mathcal{B}} - (\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_F) \frac{-\lambda'_F \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \right) \\
&= -(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_F) d \left(\frac{-\lambda'_F \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \right) - \hat{\mathcal{B}}(d\lambda_F) \frac{\lambda'_F \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \\
&= -(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_F) \left(\frac{2\lambda'_F \hat{\mathcal{Q}}^{-1} (d\lambda_F) \lambda'_F \hat{\mathcal{Q}}^{-1}}{(1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F)^2} - \frac{(d\lambda_F)' \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \right) - \hat{\mathcal{B}}(d\lambda_F) \frac{\lambda'_F \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \quad (77)
\end{aligned}$$

$$\begin{aligned}
&d\text{vec}\tilde{\mathcal{B}} \\
&= \text{vec} \left\{ -(\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_F) \left(\frac{2\lambda'_F \hat{\mathcal{Q}}^{-1} (d\lambda_F) \lambda'_F \hat{\mathcal{Q}}^{-1}}{(1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F)^2} - \frac{(d\lambda_F)' \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \right) - \hat{\mathcal{B}}(d\lambda_F) \frac{\lambda'_F \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \right\} \\
&= \frac{1}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \left(-\frac{2\hat{\mathcal{Q}}^{-1} \lambda_F \otimes (\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_F) \lambda'_F \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} + \hat{\mathcal{Q}}^{-1} \otimes (\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_F) - \hat{\mathcal{Q}}^{-1} \lambda_F \otimes \hat{\mathcal{B}} \right) d\lambda_F \\
&= \frac{1}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \left(-\frac{\hat{\mathcal{Q}}^{-1} \lambda_F \otimes (\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_F) \lambda'_F \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} + \hat{\mathcal{Q}}^{-1} \otimes (\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_F) - \hat{\mathcal{Q}}^{-1} \lambda_F \otimes \hat{\mathcal{B}} \right) d\lambda_F \\
&= \frac{1}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \left(\hat{\mathcal{Q}}^{-1} \otimes (\bar{\mathcal{R}} - \hat{\mathcal{B}}\lambda_F) + \hat{\mathcal{Q}}^{-1} \lambda_F \otimes (\hat{\mathcal{B}} - 2\tilde{\mathcal{B}}) \right) d\lambda_F \quad (78)
\end{aligned}$$

$$\begin{aligned}
&d(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}})^{-1} = -(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}})^{-1} (d(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}})) (\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}})^{-1} \\
&= -(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}})^{-1} \left((d\tilde{\mathcal{B}}') \hat{\Sigma}^{-1} \tilde{\mathcal{B}} + \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} (d\tilde{\mathcal{B}}) \right) (\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}})^{-1} \quad (79)
\end{aligned}$$

$$\begin{aligned}
& d\text{vec} \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \\
&= - \left(\left(\left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \right) \mathcal{K}_{(N-1)K} + \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \right) d\text{vec} \tilde{\mathcal{B}} \\
&= - \left(\left(\left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \right) \mathcal{K}_{(N-1)K} + \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \right) \times \\
&\quad \frac{1}{1 + \lambda_F' \hat{\mathcal{Q}}^{-1} \lambda_F} \left(- \frac{\hat{\mathcal{Q}}^{-1} \lambda_F \otimes (\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F) \lambda_F' \hat{\mathcal{Q}}^{-1}}{1 + \lambda_F' \hat{\mathcal{Q}}^{-1} \lambda_F} + \hat{\mathcal{Q}}^{-1} \otimes (\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F) - \hat{\mathcal{Q}}^{-1} \lambda_F \otimes \tilde{\mathcal{B}} \right) d\lambda_F \\
&= - \left(\left(\left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \right) \mathcal{K}_{(N-1)K} + \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \right) \times \\
&\quad \frac{1}{1 + \lambda_F' \hat{\mathcal{Q}}^{-1} \lambda_F} \left(\hat{\mathcal{Q}}^{-1} \otimes (\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F) + \hat{\mathcal{Q}}^{-1} \lambda_F \otimes (\hat{\mathcal{B}} - 2\tilde{\mathcal{B}}) \right) d\lambda_F \\
&= - (\mathcal{K}_{K^2} + I_{K^2}) \left(\left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \right) \times \\
&\quad \frac{1}{1 + \lambda_F' \hat{\mathcal{Q}}^{-1} \lambda_F} \left(\hat{\mathcal{Q}}^{-1} \otimes (\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F) + \hat{\mathcal{Q}}^{-1} \lambda_F \otimes (\hat{\mathcal{B}} - 2\tilde{\mathcal{B}}) \right) d\lambda_F \\
&= - \left\{ \frac{1}{1 + \lambda_F' \hat{\mathcal{Q}}^{-1} \lambda_F} \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} (\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F) \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \hat{\mathcal{Q}}^{-1} \right. \\
&\quad + \frac{1}{1 + \lambda_F' \hat{\mathcal{Q}}^{-1} \lambda_F} \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} (\hat{\mathcal{B}} - 2\tilde{\mathcal{B}}) \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \hat{\mathcal{Q}}^{-1} \lambda_F \\
&\quad + \frac{1}{1 + \lambda_F' \hat{\mathcal{Q}}^{-1} \lambda_F} \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \hat{\mathcal{Q}}^{-1} \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} (\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F) \\
&\quad \left. + \frac{1}{1 + \lambda_F' \hat{\mathcal{Q}}^{-1} \lambda_F} \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \hat{\mathcal{Q}}^{-1} \lambda_F \otimes \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} (\hat{\mathcal{B}} - 2\tilde{\mathcal{B}}) \right\} d\lambda_F \tag{80}
\end{aligned}$$

Therefore, we know

$$\begin{aligned}
\frac{\partial \text{GLS-LM}}{\partial \lambda'_F} &= \frac{\partial FAR}{\partial \lambda'_F} + \frac{2T}{\left(1 + \lambda'_F \widehat{Q}^{-1} \lambda_F\right)^2} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}}\right)^{-1} \widehat{Q}^{-1} \times \\
&\quad \left(\left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right)' \widehat{\Sigma}^{-1/2} \mathcal{M}_{\widehat{\Sigma}^{-1/2} \widetilde{\mathcal{B}}} \widehat{\Sigma}^{-1/2} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right) + \lambda_F \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right)' \widehat{\Sigma}^{-1/2} \mathcal{M}_{\widehat{\Sigma}^{-1/2} \widetilde{\mathcal{B}}} \widehat{\Sigma}^{-1/2} \left(\widehat{\mathcal{B}} - 2\widetilde{\mathcal{B}}\right) \right) \\
&= \frac{\partial FAR}{\partial \lambda'_F} + \frac{2T}{\left(1 + \lambda'_F \widehat{Q}^{-1} \lambda_F\right)^2} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \times \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}}\right)^{-1} \widehat{Q}^{-1} \times \\
&\quad \left\{ \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right)' \widehat{\Sigma}^{-1/2} \mathcal{M}_{\widehat{\Sigma}^{-1/2} \widetilde{\mathcal{B}}} \widehat{\Sigma}^{-1/2} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right) \right. \\
&\quad \left. - \lambda_F \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right)' \widehat{\Sigma}^{-1/2} \mathcal{M}_{\widehat{\Sigma}^{-1/2} \widetilde{\mathcal{B}}} \widehat{\Sigma}^{-1/2} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right) \lambda'_F \frac{\widehat{Q}^{-1}}{1 + \lambda'_F \widehat{Q}^{-1} \lambda_F} \right\} \\
&= \frac{\partial FAR}{\partial \lambda'_F} - \frac{\partial FAR}{\partial \lambda'_F} \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}}\right)^{-1} \left(\widehat{Q}^{-1} - \frac{\widehat{Q}^{-1} \lambda_F \lambda'_F \widehat{Q}^{-1}}{1 + \lambda'_F \widehat{Q}^{-1} \lambda_F} \right) \times \\
&\quad \frac{\left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right)' \widehat{\Sigma}^{-1/2} \mathcal{M}_{\widehat{\Sigma}^{-1/2} \widetilde{\mathcal{B}}} \widehat{\Sigma}^{-1/2} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F\right)}{1 + \lambda'_F \widehat{Q}^{-1} \lambda_F} \tag{83}
\end{aligned}$$

Therefore, we have the results in proposition 4.

Q.E.D.

Proposition 5.

$$\frac{\partial JGLS}{\partial \lambda'_F} = \frac{JGLS(\lambda_F)}{T} \frac{\partial FAR}{\partial \lambda'_F} \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}}\right)^{-1} \mathcal{Q}_- \tag{84}$$

Proof of proposition 5: This is a direct result of $JGLS = FAR - GLS-LM$ such that

$$\begin{aligned}
\frac{\partial JGLS}{\partial \lambda'_F} &= \frac{\partial FAR}{\partial \lambda'_F} - \frac{\partial GLS-LM}{\partial \lambda'_F} \\
&= \frac{\partial FAR}{\partial \lambda'_F} \left(\tilde{\mathbf{B}}' \hat{\Sigma}^{-1} \tilde{\mathbf{B}} \right)^{-1} \left(\hat{\mathbf{Q}}^{-1} - \frac{\hat{\mathbf{Q}}^{-1} \lambda'_F \lambda'_F \hat{\mathbf{Q}}^{-1}}{1 + \lambda'_F \hat{\mathbf{Q}}^{-1} \lambda'_F} \right) \times \\
&\quad \frac{\left(\bar{\mathbf{R}} - \hat{\mathbf{B}} \lambda'_F \right)' \hat{\Sigma}^{-1/2} \mathcal{M}_{\hat{\Sigma}^{-1/2} \tilde{\mathbf{B}}} \hat{\Sigma}^{-1/2} \left(\bar{\mathbf{R}} - \hat{\mathbf{B}} \lambda'_F \right)}{1 + \lambda'_F \hat{\mathbf{Q}}^{-1} \lambda'_F} \\
&= \frac{JGLS(\lambda'_F)}{T} \frac{\partial FAR}{\partial \lambda'_F} \left(\tilde{\mathbf{B}}' \hat{\Sigma}^{-1} \tilde{\mathbf{B}} \right)^{-1} \left(\hat{\mathbf{Q}}^{-1} - \frac{\hat{\mathbf{Q}}^{-1} \lambda'_F \lambda'_F \hat{\mathbf{Q}}^{-1}}{1 + \lambda'_F \hat{\mathbf{Q}}^{-1} \lambda'_F} \right) \quad (85)
\end{aligned}$$

Q.E.D.

Proposition 6.

$$\begin{aligned}
&\frac{\partial^2 FAR}{\partial \lambda'_F \partial \lambda'_F} \\
&= \left\{ \left(\frac{2T}{1 + \lambda'_F \hat{\mathbf{Q}}^{-1} \lambda'_F} \tilde{\mathbf{B}}' \hat{\Sigma}^{-1} \tilde{\mathbf{B}} \right) + \frac{2T}{\left(1 + \lambda'_F \hat{\mathbf{Q}}^{-1} \lambda'_F \right)^2} \times \right. \\
&\quad \left. \left\{ \tilde{\mathbf{B}}' \hat{\Sigma}^{-1} \left(\bar{\mathbf{R}} - \hat{\mathbf{B}} \lambda'_F \right) \lambda'_F \hat{\mathbf{Q}}^{-1} + \hat{\mathbf{Q}}^{-1} \lambda'_F \left(\bar{\mathbf{R}} - \hat{\mathbf{B}} \lambda'_F \right)' \hat{\Sigma}^{-1} \tilde{\mathbf{B}} + \left(\bar{\mathbf{R}} - \hat{\mathbf{B}} \lambda'_F \right)' \hat{\Sigma}^{-1} \left(\bar{\mathbf{R}} - \hat{\mathbf{B}} \lambda'_F \right) \times \right. \right. \\
&\quad \left. \left. \left(\frac{\hat{\mathbf{Q}}^{-1} \lambda'_F \otimes \lambda'_F \hat{\mathbf{Q}}^{-1}}{1 + \lambda'_F \hat{\mathbf{Q}}^{-1} \lambda'_F} - \hat{\mathbf{Q}}^{-1} \right) \right\} \right\} \quad (86)
\end{aligned}$$

Proof of proposition 6:

$$\begin{aligned}
& d\left(\frac{\partial FAR}{\partial \lambda'_F}\right)' = \text{dvec}\left(\frac{\partial FAR}{\partial \lambda'_F}\right)' = d\left(\frac{\partial FAR}{\partial \lambda_F}\right) = d\left(-\frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F)\right) \\
& = \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F) \frac{4T \lambda'_F \widehat{\mathcal{Q}}^{-1} d\lambda_F}{(1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} - \frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} (d\widetilde{\mathcal{B}})' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F) \\
& \quad + \frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widehat{\mathcal{B}} d\lambda_F \\
& = (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \frac{4T \lambda'_F \widehat{\mathcal{Q}}^{-1}}{(1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} d\lambda_F \\
& \quad - \frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \left((\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F)' \widehat{\Sigma}^{-1} \otimes I_K \right) \mathcal{K}_{(N-1)K} (d\text{vec} \widetilde{\mathcal{B}}) \\
& \quad + \left(\frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widehat{\mathcal{B}} \right) d\lambda_F \\
& = \left\{ \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F) \frac{4T \lambda'_F \widehat{\mathcal{Q}}^{-1}}{(1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} \right) \right. \\
& \quad - \frac{2T}{(1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} \left((\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F)' \widehat{\Sigma}^{-1} \otimes I_K \right) \mathcal{K}_{(N-1)K} \times \\
& \quad \left(-\frac{\widehat{\mathcal{Q}}^{-1} \lambda_F \otimes (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F) \lambda'_F \widehat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} + \widehat{\mathcal{Q}}^{-1} \otimes (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F) - \widehat{\mathcal{Q}}^{-1} \lambda_F \otimes \widetilde{\mathcal{B}} \right) \\
& \quad \left. + \left(\frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widehat{\mathcal{B}} \right) \right\} d\lambda_F \\
& = \left\{ \left(\frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widehat{\mathcal{B}} \right) + \frac{2T}{(1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} \times \right. \\
& \quad \left\{ \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F) \lambda'_F \widehat{\mathcal{Q}}^{-1} + \widehat{\mathcal{Q}}^{-1} \lambda_F (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} + (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F)' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F) \times \right. \\
& \quad \left. \left. \left(\frac{\widehat{\mathcal{Q}}^{-1} \lambda_F \otimes \lambda'_F \widehat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} - \widehat{\mathcal{Q}}^{-1} \right) \right\} \right\} d\lambda_F \tag{87}
\end{aligned}$$

Therefore, we know the Hessian matrix of FAR statistic evaluated at the zero point of $\frac{\partial FAR}{\partial \lambda_F}$, λ_F^* , equals

$$\frac{\partial^2 FAR}{\partial \lambda_F \partial \lambda_F'} \Big|_{\lambda_F^*} = \frac{2T}{1 + \lambda_F^* \widehat{\mathcal{Q}}^{-1} \lambda_F^*} \left\{ \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} - \frac{FAR(\lambda_F^*)}{T} \mathcal{Q}_- \right\} \quad (88)$$

where $\mathcal{Q}_- = \left(\widehat{\mathcal{Q}}^{-1} - \frac{\widehat{\mathcal{Q}}^{-1} \lambda_F \otimes \lambda_F' \widehat{\mathcal{Q}}^{-1}}{1 + \lambda_F' \widehat{\mathcal{Q}}^{-1} \lambda_F} \right)$ and \mathcal{Q}_- is positive definite.

Q.E.D.

Proposition 7.

$$\begin{aligned} \frac{\partial^2 JGLS}{\partial \lambda_F \partial \lambda_F'} &= JGLS(\lambda_F)/T \times \left\{ \frac{2T}{1 + \lambda_F' \widehat{\mathcal{Q}}^{-1} \lambda_F} \mathcal{Q}_- \right. \\ &+ \frac{2T \left[\mathcal{Q}_- \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right) \lambda_F \widehat{\mathcal{Q}}^{-1} + \widehat{\mathcal{Q}}^{-1} \lambda_F \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \mathcal{Q}_- \right]}{(1 + \lambda_F' \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} \\ &- \frac{2T}{1 + \lambda_F' \widehat{\mathcal{Q}}^{-1} \lambda_F} FAR(\lambda_F) \mathcal{Q}_- \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \mathcal{Q}_- \\ &+ \frac{2}{1 + \lambda_F' \widehat{\mathcal{Q}}^{-1} \lambda_F} GLS-LM(\lambda_F) \mathcal{Q}_- \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \mathcal{Q}_- \\ &+ \left. \frac{2T}{(1 + \lambda_F' \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} \mathcal{Q}_- \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right) \left(\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F \right)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \mathcal{Q}_- \right\} \\ &+ JGLS(\lambda_F)/T^2 \mathcal{Q}_- \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \frac{\partial FAR}{\partial \lambda_F} \frac{\partial FAR}{\partial \lambda_F'} \left(\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} \right)^{-1} \mathcal{Q}_- \end{aligned} \quad (89)$$

Proof of proposition 7:

$$\begin{aligned}
& d\left(\frac{\partial JGLS}{\partial \lambda_F}\right) \\
&= JGLS(\lambda_F)/T \left(\hat{Q}^{-1} - \frac{\hat{Q}^{-1} \lambda_F \lambda'_F \hat{Q}^{-1}}{1 + \lambda'_F \hat{Q}^{-1} \lambda_F} \right) (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} d\left(\frac{\partial FAR}{\partial \lambda'_F}\right)' \\
&\quad + JGLS(\lambda_F)/T d\left(\left(\hat{Q}^{-1} - \frac{\hat{Q}^{-1} \lambda_F \lambda'_F \hat{Q}^{-1}}{1 + \lambda'_F \hat{Q}^{-1} \lambda_F} \right) (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} \right) \frac{\partial FAR}{\partial \lambda_F} \\
&\quad + d(JGLS(\lambda_F)/T) \left(\hat{Q}^{-1} - \frac{\hat{Q}^{-1} \lambda_F \lambda'_F \hat{Q}^{-1}}{1 + \lambda'_F \hat{Q}^{-1} \lambda_F} \right) (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} \frac{\partial FAR}{\partial \lambda_F} \\
&= JGLS(\lambda_F)/T \left(\hat{Q}^{-1} - \frac{\hat{Q}^{-1} \lambda_F \lambda'_F \hat{Q}^{-1}}{1 + \lambda'_F \hat{Q}^{-1} \lambda_F} \right) (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} d\left(\frac{\partial FAR}{\partial \lambda'_F}\right)' \\
&\quad + JGLS(\lambda_F)/T d\left(\left(\hat{Q}^{-1} - \frac{\hat{Q}^{-1} \lambda_F \lambda'_F \hat{Q}^{-1}}{1 + \lambda'_F \hat{Q}^{-1} \lambda_F} \right) (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} \right) \frac{\partial FAR}{\partial \lambda_F} \\
&\quad + JGLS(\lambda_F)/T^2 \frac{\partial FAR}{\partial \lambda'_F} (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} \left(\hat{Q}^{-1} - \frac{\hat{Q}^{-1} \lambda_F \lambda'_F \hat{Q}^{-1}}{1 + \lambda'_F \hat{Q}^{-1} \lambda_F} \right) (d\lambda_F) \times \\
&\quad \left(\hat{Q}^{-1} - \frac{\hat{Q}^{-1} \lambda_F \lambda'_F \hat{Q}^{-1}}{1 + \lambda'_F \hat{Q}^{-1} \lambda_F} \right) (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} \frac{\partial FAR}{\partial \lambda_F} \tag{90}
\end{aligned}$$

$$\begin{aligned}
& d\left(\left(\hat{Q}^{-1} - \frac{\hat{Q}^{-1} \lambda_F \lambda'_F \hat{Q}^{-1}}{1 + \lambda'_F \hat{Q}^{-1} \lambda_F} \right) (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} \right) \frac{\partial FAR}{\partial \lambda_F} \\
&= \left\{ \frac{2T}{(1 + \lambda'_F \hat{Q}^{-1} \lambda_F)^2} Q_- (\bar{R} - \hat{B} \lambda_F)' \hat{\Sigma}^{-1} \tilde{B} (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} \hat{Q}^{-1} \lambda_F \right. \\
&\quad + \frac{2T}{(1 + \lambda'_F \hat{Q}^{-1} \lambda_F)^2} \hat{Q}^{-1} \lambda_F (\bar{R} - \hat{B} \lambda_F)' \hat{\Sigma}^{-1} \tilde{B} (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} Q_- \\
&\quad + \frac{2}{1 + \lambda'_F \hat{Q}^{-1} \lambda_F} GLS-LM(\lambda_F) Q_- (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} Q_- \\
&\quad - \frac{2T}{(1 + \lambda'_F \hat{Q}^{-1} \lambda_F)^2} Q_- (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} \hat{Q}^{-1} \lambda_F (\bar{R} - \hat{B} \lambda_F)' \hat{\Sigma}^{-1} \tilde{B} \\
&\quad + \frac{2T}{(1 + \lambda'_F \hat{Q}^{-1} \lambda_F)^2} Q_- (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} \tilde{B}' \hat{\Sigma}^{-1} (\bar{R} - \hat{B} \lambda_F) (\bar{R} - \hat{B} \lambda_F)' \hat{\Sigma}^{-1} \tilde{B} (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} Q_- \\
&\quad \left. - \frac{2T}{(1 + \lambda'_F \hat{Q}^{-1} \lambda_F)^2} Q_- (\bar{R} - \hat{B} \lambda_F)' \hat{\Sigma}^{-1} \tilde{B} (\tilde{B}' \hat{\Sigma}^{-1} \tilde{B})^{-1} \hat{Q}^{-1} \lambda_F \right\} d\lambda_F \tag{91}
\end{aligned}$$

$$\begin{aligned}
& \left(\hat{\mathcal{Q}}^{-1} - \frac{\hat{\mathcal{Q}}^{-1} \lambda_F \lambda'_F \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \right) \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} d \left(\frac{\partial FAR}{\partial \lambda'_F} \right)' \\
&= \left\{ \frac{2T}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \mathcal{Q}_- \right. \\
&\quad + \frac{2T}{(1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F)^2} \mathcal{Q}_- \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right) \lambda_F \hat{\mathcal{Q}}^{-1} \\
&\quad + \frac{2T}{(1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F)^2} \mathcal{Q}_- \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \hat{\mathcal{Q}}^{-1} \lambda_F \left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right)' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \\
&\quad \left. - \frac{2T}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} FAR(\lambda_F) \mathcal{Q}_- \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \mathcal{Q}_- \right\} d\lambda_F \tag{92}
\end{aligned}$$

$$\begin{aligned}
& d \left(\left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right)' \hat{\Sigma}^{-1/2} \mathcal{M}_{\hat{\Sigma}^{-1/2} \tilde{\mathcal{B}}} \hat{\Sigma}^{-1/2} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right) \right) \\
&= 2 \left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right)' \hat{\Sigma}^{-1/2} \mathcal{M}_{\hat{\Sigma}^{-1/2} \tilde{\mathcal{B}}} \hat{\Sigma}^{-1/2} d\lambda_F \\
&\quad - \left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right)' \hat{\Sigma}^{-1} d \left(\tilde{\mathcal{B}} \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \tilde{\mathcal{B}}' \right) \hat{\Sigma}^{-1} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right) \\
&= 2 \left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right)' \hat{\Sigma}^{-1/2} \mathcal{M}_{\hat{\Sigma}^{-1/2} \tilde{\mathcal{B}}} \hat{\Sigma}^{-1/2} d\lambda_F \\
&\quad - 2 \frac{\left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right)' \hat{\Sigma}^{-1/2} \mathcal{M}_{\hat{\Sigma}^{-1/2} \tilde{\mathcal{B}}} \hat{\Sigma}^{-1/2} \left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right)}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \times \\
&\quad \left(\bar{\mathcal{R}} - \hat{\mathcal{B}} \lambda_F \right)' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \left(\tilde{\mathcal{B}}' \hat{\Sigma}^{-1} \tilde{\mathcal{B}} \right)^{-1} \left(\hat{\mathcal{Q}}^{-1} - \frac{\hat{\mathcal{Q}}^{-1} \lambda_F \lambda'_F \hat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \hat{\mathcal{Q}}^{-1} \lambda_F} \right) d\lambda_F \tag{93}
\end{aligned}$$

$$\begin{aligned}
& d \left(\left(\widehat{\mathcal{Q}}^{-1} - \frac{\widehat{\mathcal{Q}}^{-1} \lambda_F \lambda'_F \widehat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \right) (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \right) \\
&= \left(\frac{\widehat{\mathcal{Q}}^{-1} \lambda_F 2 \lambda'_F \widehat{\mathcal{Q}}^{-1} (d \lambda_F) \lambda'_F \widehat{\mathcal{Q}}^{-1}}{(1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} - \frac{\widehat{\mathcal{Q}}^{-1} (d \lambda_F) \lambda'_F \widehat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} - \frac{\widehat{\mathcal{Q}}^{-1} \lambda_F (d \lambda'_F) \widehat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \right) (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \\
&+ \left(\widehat{\mathcal{Q}}^{-1} - \frac{\widehat{\mathcal{Q}}^{-1} \lambda_F \lambda'_F \widehat{\mathcal{Q}}^{-1}}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \right) d (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \tag{94}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial^2 JGLS}{\partial \lambda_F \partial \lambda'_F} &= JGLS(\lambda_F)/T \times \left\{ \frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} \mathcal{Q}_- \right. \\
&+ \frac{2T \left[\mathcal{Q}_- (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F) \lambda_F \widehat{\mathcal{Q}}^{-1} + \widehat{\mathcal{Q}}^{-1} \lambda_F (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \mathcal{Q}_- \right]}{(1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} \\
&- \frac{2T}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} FAR(\lambda_F) \mathcal{Q}_- (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \mathcal{Q}_- \\
&+ \frac{2}{1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F} GLS-LM(\lambda_F) \mathcal{Q}_- (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \mathcal{Q}_- \\
&+ \left. \frac{2T}{(1 + \lambda'_F \widehat{\mathcal{Q}}^{-1} \lambda_F)^2} \mathcal{Q}_- (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F) (\bar{\mathcal{R}} - \widehat{\mathcal{B}} \lambda_F)' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}} (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \mathcal{Q}_- \right\} \\
&+ JGLS(\lambda_F)/T^2 \mathcal{Q}_- (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \frac{\partial FAR}{\partial \lambda_F} \frac{\partial FAR}{\partial \lambda'_F} (\widetilde{\mathcal{B}}' \widehat{\Sigma}^{-1} \widetilde{\mathcal{B}})^{-1} \mathcal{Q}_- \tag{95}
\end{aligned}$$

Q.E.D.